

Changing universe model with applications

John C. Hodge*

Blue Ridge Community College, 100 College Dr., Flat Rock, NC, 28731-1690

(Dated: February 2, 2008)

Although the current galaxy models yield calculations consistent with much of the data, many irregularities exist, exceptions have been found to the current models, the Λ CDM model apparently fails on galaxy scales, dark matter remains elusive, the phenomena at the center of galaxies are only beginning to be addressed, and many observations and empirical relationships are unexplained. A changing universe model (CUM) is proposed that posits the stuff of our universe is continually erupting into our universe from sources at the center of galaxies. In this first test of the CUM, a single equation that describes the rotation velocity curve of spiral galaxies is derived. The equation is also used to correlate the measured mass of the theorized, central supermassive black hole with other galaxy parameters. The equation adds one parameter for each mature galaxy and one parameter for each particle (atom nucleus and smaller) species to Newtonian dynamics. Rising, flat, and declining rotation velocity curves are explained without unknown dark matter. Ten previously unknown parametric relationships are discovered. Also, the CUM integrates several heretofore unrelated observations. The CUM model is in an early development stage and has already predicted new relationships among galaxy parameters.

PACS numbers: 98.80.Bp, 98.62.Ai, 98.62.Dm, 98.62.Js

I. INTRODUCTION

The discrepancy between the Newtonian estimated mass in galaxies and galaxy clusters and the observation of star and gas kinematics is well established. Currently, the possible explanations for this discrepancy are that a large mass M_{DM} of dark matter (DM) and dark energy exists (the Λ CDM model) or that Newtonian gravity is modified on galactic scales. The current Λ CDM model of the structure of the universe appears successful on large scales [1], but galactic scale predictions disagree with observations [2, 3, 4]. The most popular modified gravity model is proposed by the Modified Newtonian Dynamics (MOND) model [5, and references therein]. MOND suggests gravitational acceleration changes with galactic distance scales. MOND appears limited to only the disk region of spiral galaxies and appears to have two falsifiers [5]. How MOND may be applied to cosmological scales is unclear. However, MOND may represent an *effective* force law arising from a broader force law [6]. The evidence suggest the problem of a single model explaining both galactic scale and cosmological scale observations is fundamental [4].

An alternate explanation is the possibility there exists another force originating from the center of galaxies. This proposal, called the changing universe model (CUM), posits the stuff of our universe is matter and a matterless scalar ρ whose gradient exerts a force $\mathbf{F}_s$ on the cross section of matter. The $\mathbf{F}_s$ is outward from galactic center. Thus, $\mathbf{F}_s$ is a repulsive force in a galaxy relative to gravity. However, it differs from a gravity type force since it acts on the cross section of matter and it is

matterless.

The rotation velocity v (km s^{-1}) of a particle in the plane of a spiral galaxy (hereinafter galaxy) reflects the forces acting on the particle. An explanation of v^2 as a function of radius r (kpc) from the center of a galaxy (RC) requires a knowledge of galaxy dynamics and an explanation of radically differing slopes. The v and, thus, the RC is measured along the major axis. Although v is non-relativistic, the result of calculations using RC models are used in cosmological calculations. Thus, the RC is an excellent tool to evaluate galaxy models. Further, a complete explanation of the RC must also explain black holes and observations of the center of galaxies.

Battaner and Florido [7] and Sofue and Rubin [8] provides an overview of the current state of knowledge of RC's in the outer bulge and disk regions of galaxies. The particles most often measured in the disk region of a galaxy are hydrogen gas by H I observation and stars by observing the H_α line. The particles being measured in the inner bulge region are stars by observation of the H_α , CO, and other spectral lines. Also, the RC differs for different particles. For example, although the H I and H_α RCs for NGC4321 [9] in Fig. 1 differ in the outer bulge, they approach each other in the outer disk region. Most models of RC monitor the H I in the outer bulge and disk regions and ignore the differing H_α RC. Also, when drawing the H I RC, a straight line is often drawn through the center. This practice is inconsistent with stellar observations [10].

The observation of rising RCs in the outer bulge and rising, flat, or declining RC's in the disk region of galaxies poses a perplexing problem. If most of the matter of a galaxy is in the bulge region, classical Newtonian mechanics predicts a Keplerian declining RC in the disk.

Currently, a convention on how to calculate error (e.g. random or systematic) in RCs is lacking in the literature.

*jch9496@blueridge.edu




FIG. 1: Plot of the H I (filled diamonds) and H α (filled squares) RC for NGC4321.

One popular method is based on the assumption of symmetry. This practice of averaging the v of the approaching and receding sides of the RC is ignoring important information [11]. Asymmetry appears to be the norm rather than the exception [12]. The RC of all nearby galaxies for which the kinematics have been studied have asymmetry [12]. At larger distance, objects appear to become more symmetric as they become less well-resolved (H I resolution is typically 15 arcsec to 50 arcsec) [11].

Hodge and Castelaz [13] found a correlation between the size and distance of neighboring galaxies and the asymmetry in the disk of the H I RC. The galaxies studied with minimal asymmetry also had rising RCs. Therefore, rising RCs are intrinsic to galaxies.

A strong correlation between the core radius R_c and the stellar exponential scale length R_d was found by Donato and Salucci [14]. The R_d is derived from photometric measurements. The R_c is obtained from kinematic measurements. If this relationship holds for both high surface brightness (HSB) galaxies, low surface brightness (LSB) galaxies, galaxies with rising RCs, and galaxies with declining RCs, modeling such a relationship is very difficult using current galaxy models.

Ghez et al. [15] and Ferrarese and Merritt [16] have observed Keplerian motion to within one part in 100 in elliptical orbits of stars that are from less than a pc to a few 1000 pc from the center of galaxies.

The orbits of stars within nine light hours of the Galaxy center indicates the presence of a large amount of mass within the orbits [10, 15, 17, 18, 19]. To achieve the velocities of 1300 km s^{-1} to 9000 km s^{-1} [10] and high accelerations [15], there must be a huge amount of very dense particles such as millions of black holes, dense quark stars [20, and references therein], and ionized iron [21] distributed inside the innermost orbit of luminous matter. The mass $M_\bullet$ (in $M_\odot$) within a few light

hours of the center of galaxies varies from $10^6 M_\odot$ to $10^{10} M_\odot$ [22, 23]. The $M_\bullet$ can be distributed over the volume with a density of at least $10^{12} M_\odot \text{ pc}^{-3}$ [15, 18, 19, 24]. The orbits of stars closest to the center are approximately 1,169 times [10] the Schwarzschild radius of the theorized supermassive black hole (SBH) [15]. However, such large mass crowded into a ball with a radius of less than 60 AU must either quickly dissipate, must quickly collapse into a SBH [25, 26], or there must exist a force in addition to the centrifugal force holding the mass from collapse. Mouawad et al. [27] suggested there is some extended mass around Sgr A. Also, models of supermassive fermion balls wherein the gravitational pressure is balanced by the degeneracy pressure of the fermions due to the Pauli exclusion principle are not excluded [28]. A strong repulsive force at the center of galaxies would explain the shells of shocked gas around galactic nuclei [29, page 595][30], the apparent inactivity of the central object [31, 32, 33, 34], and the surprising accuracy of reverberation mapping as a technique to determine $M_\bullet$ [35]. However, discrepancies have been noted among the methods used to measure $M_\bullet$ [35, 36, and references therein].

The first published calculation of the slope of the $M_\bullet$ to velocity dispersion σ_v (in km s^{-1}) curve ($M_\bullet - \sigma_v$ relation) varied between 5.27 ± 0.40 [37] and 3.75 ± 0.3 [23]. Reference [38, and references therein] suggested the range of slopes are caused by systematic differences in the velocity dispersions used by different groups. However, the origin of these differences remains unclear.

Ferrarese [39] found the ratio of the $M_\bullet$ to M_{DM} of the theorized dark matter halo around galaxies was a positive value that decreased with halo mass. However, if the intrinsic rotation curve is rising [13], the effect of the force of M_{DM} in the equations implies the effect of the center object must be repulsive. Such a repulsive force was called a “wind” by Shu et al. [40], Silk and Rees [41]. The “wind” (a gas) exerted a repulsive force acting on the cross sectional area of particles. Therefore, denser particles such as black holes move inward relative to less dense particles.

A multitude of X-ray point sources, highly ionized iron, and radio flares without accompanying large variation at longer wavelengths have been reported near the center of the Milky Way [21, 29, 31, 32, 34, 42, 43].

McLure and Dunlop [44] found that the mass in the central region is 0.0012 of the mass of the bulge. Ferrarese and Merritt [16] reported that approximately 0.1% of a galaxy’s luminous mass is at the center of galaxies and that the density of SBH’s in the universe agrees with the density inferred from observation of quasars. Merritt and Ferrarese [45] found similar results in their study of the $M_\bullet - \sigma_v$ relation. Ferrarese [39] found a tight relation between rotation velocity v_c in the outer disk region and bulge velocity dispersion σ_c ($v_c - \sigma_c$) which is strongly supporting a relationship of a center force with total gravitational mass of a galaxy. Wandel [46] showed the $M_\bullet$ of AGN galaxies and their bulge luminosity follow the same relationships as their ordinary (inactive) galaxies,

with the exception of narrow line AGN. Graham et al. [47, 48, 49] found correlations between $M_\bullet$ and structural parameters of elliptical galaxies and bulges. Either the dynamics of many galaxies are producing the same increase of mass in the center at the same rate or a feedback controlled mechanism exists to evaporate the mass increase that changes as the rate of inflow changes as suggested by Merritt and Ferrarese [35]. The RCs imply the dynamics of galaxies differ so the former explanation is unlikely.

In accordance with the Principle of Fundamental Principles (see Appendix A), a search was made for a physical process in the Newtonian physical domain that may model and explain the observed data of spiral galaxies. Such a system was found in the fractional distillation process (FDP). A FDP has a Source of heat (Space energy) and material input (matter) at the bottom (center of galaxies) of a container. The heat of the FDP is identified with a constituent of our universe that is not matter and is identified as Space (in units of “que”, abbreviated as “q”, - a new unit of measure) whose density ρ is a potential energy. The term “Space” will mean the amount of Space in ques. In the FDP, temperature decreases with height from the Source. The opposing forces of gravity and the upward flow of gas acting on the cross section of molecules in an FDP cause the compounds to be lighter with height and lower temperature. Heavy molecules that coalesce and fall to the bottom are re-evaporated and cracked. The heat equation with slight modification rather than gas dynamics modeled the flow of heat (Space energy). Temperature decline of the FDP was modeled as dissipation of matter and Space energy into three dimensions. Thus, the Changing Universe Model (CUM) was born.

This paper posits that there is a Source of strength ϵ at the center of each galaxy. The Source is a source of all matter and of Space. The ρ forms a scalar potential field. Energy is conserved in the emission of matter/energy and Space from the Source. The unit of measure of ϵ is in the amount of que released per second which is proportional to the amount of mass released per second, ϵ_m in units of $M_\odot \text{ s}^{-1}$. The application of the CUM to the RC of galaxies explains observations and empirical relations. Also, ten new relationships among galaxy parameters were predicted and confirmed. This, the first test of the CUM, successfully explains RC observations and suggests an alternative to DM. The CUM, at this early stage of model development, is bringing together heretofore apparently unrelated observations.

The object of this article is to develop and test the CUM by suggesting a galaxy model consistent with the CUM, by explaining observations of the RC and the centers of galaxies, and by deriving relations among galaxy parameters. In section II the CUM model is developed from the fundamental principles, which are stated in Appendix A. An equation for the RC is developed from the energy equations of the CUM, applied to galaxies, and compared with galaxy observations in Section III.

The posited description of particles is used to explain several phenomena involving observations of the center of galaxies. Other parameter relationships are presented in Section IV. Section V shows the interger relationships developed in the previous sections are not random. The results are discussed in Section VI. Section VII lists the conclusions.

II. CUM MODEL

The assumed Fundamental Principles are stated in Appendix A. Throughout the text, capitalizing the first letters of the name of the Fundamental Principle denotes these principles.

The Principles of Change and Repetition requires that the three dimensions (3D) be created from two dimensions (2D). The creation of 3D from zero dimensions (0D) is not only unlikely but forbidden. The universe begins with a Change that may be a rupture of 0D. The Change itself is a beginning of time. The 0D ruptures to create one dimensional (1D) points and Space between the points on a line. The 0D is a Source of 1D and continually ejects Space into one end of the 1D line. The other end must expand into the void. Thus, the 1D line is continually growing. The 1D line has only one direction. Therefore, the 1D line is straight. The 1D energy is $\tau_1 t = \rho_1 l$ where τ_1 is the point tension, t is time elapsed between point eruption, ρ_1 is the line Space density, and l is the length between points. The first transition creates length and time

$$l = ct, \quad (1)$$

where $c = \tau_1/\rho_1$ is the constant of proportionality.

When points in 1D rupture, the 2D Space is inserted into 2D at many points along the 1D line. The rupture can have no component remaining in 1D. Thus, a right angle is defined and the metric coefficients equal one (a Cartesian plane). Accordingly, 2D is flat. The 2D energy for each transition is $\tau_2 l = \rho_2 S$ where τ_2 is the line tension, l is the line length, ρ_2 is the area Space density, and S is the area.

A 2D Sink is at a place in 2D that has Space tightly contained in a line. The line is a one-dimensional border in 2D. If a line with a “line tension” compressing Space is deformed perpendicular to the line, an arc of height h is formed. The geometric constant of the line is its length l . If a condition of $l = \pi h$ is reached, a bubble in 2D is formed. The end points of the line are coincident. This rupture causes Space to be removed from the 2D plane. This 2D Sink becomes a Source in 3D Space. The line becomes an object in 3D called a hod. The hod is the smallest piece of matter from which all other matter is constructed. The hod is supple so that its shape may change in 3D. The edge of the 2D plane expands into the void. Hence, the distance between the points that rupture into 3D is increasing. The expansion is recurrent and is in 2D, not 3D. Because the rupture into 3D

is orthogonal to 2D, 3D is flat (Cartesian three space dimensions), also. The l becomes area α . The Space area in 2D becomes the volume V of Space density ρ in 3D. One que is the amount of Space released by one hod in the transition.

Since nothing exists in 3D except because of the transitions, the energy for a transition is

$$\tau\alpha = \rho_I V_I, \quad (2)$$

where ρ_I and V_I are the initial ρ and V at the transition, respectively, and τ is the proportionality constant called surface tension.

Similarly, corollary II of the Fundamental Principle implies a 3D Sink may be formed.

Succeeding ruptures at the same point adds Space and hods to the universe. Since we observe there is Space and ρ in our neighborhood, the Space and hods then flow from the Source and creates the galaxy around the Source.

The total energy transferred into 3D from Eq. (2) is

$$N_t \tau\alpha = N_t \rho_I V_I, \quad (3)$$

where N_t is the total number of hods in the universe.

In the observed universe, the 2D plane is common for all Sources. As the Space from a Source expands into the void, it encounters other Space from other Sources. Since all Space originates from one 0D Source, the merged Space is causally correlated and coherent. However, 2D Space is distinct from 3D Space.

In 2D, the surface was a tightly bound line on the outside of Space. In 3D, Space is on the outside of a surface. The τ on the 2D Space was high enough to contain the Space. Now τ acts to attract 3D Space. Since the hod formed in the extreme pressure of the Source, the τ maximum ability to act on Space exceeds the pressure of Space on the hod away from the Source. Since τ and α are constant, as the extent of the volume V increases so ρ must decrease. Therefore, the hod applies its energy, $\tau\alpha$, on the Space immediately surrounding the hod. Since the hod contained the Space before release and Space was at a much higher ρ than is surrounding the hod in 3D, all the hod's tensional energy is exerted on Space. A hod is a 2D discontinuity in the continuity of 3D Space. These discontinuities are surfaces in Space with an edge not containing loops.

Hods are discrete. The mathematics of integer numbers can then be used to count hods. In addition, hods can combine, one with another. The number of hods and the structure of the combinations of hods create the different particles and forces, except gravity, of the universe. Therefore, the language of hods is integer numbers, structures, counting, subtraction, and summation.

Space is continuous, perfectly elastic, and compressible. The Space continuously fills the region it occupies except for the hods. Transformation of Space over time is continuous. Hods and the outer edge of Space as it expands into the void are discontinuities in Space. Space really can meet the mathematical criteria of considering

an arbitrarily small volume. Therefore, the language of Space is real numbers, tensors, field theories, differentiation, and integrals.

The three interactions of concern are that each hod exerts a force on Space, Space exerts a force on the hod, and Space exerts a force on adjacent Space. A hod is not attracted or repelled by other hods through a field or other action at a distance device. Since τ can act only perpendicular to the hod's surface, the force Space exerts on the hod is perpendicular to the surface of the hod.

There are three dimensions in our universe. Counting is a characteristic of hods. Therefore, that our universe is three-dimensional is a characteristic of the hods and not of Space.

There are two concepts of distance. Distance can be by reference to a fixed grid. Traditional methods have used Space as the fixed grid. In this model, distance is a multiple of the average diameter a hod. Since α is a constant and the hod may be deformed, the hod average diameter will be considered a constant throughout the universe beyond the Source and in moving and accelerating reference frames. The measure of distance will be a "ho", which is defined as the average diameter of the hod. The CUM uses ho distance to form a grid in a flat universe with

$$1 \text{ ho} = 2 \left(\frac{\alpha}{\pi} \right)^{1/2}. \quad (4)$$

Another concept of distance is the amount of Space between two surfaces. By analogy, this type of distance in gas is the amount of gas between two surfaces. Increasing the amount of gas between the two surfaces increases the distance. One way to do this is by movement of the surfaces. Another way is to add more gas from a source. If the medium is compressible, the distance measured by the amount of gas varies with the change in density of the medium. The CUM model allows Space to be compressible. Therefore, the amount of Space between two surfaces can change if the ρ (que ho⁻³) of the Space between two surfaces changes.

The terms distance, length, area, and volume will refer to ho measurements. The unit of measure of distance is usually expressed in km, pc, kpc, or Mpc. If the measurement is taken by means of angular observation or time-between-events observations such as using Cepheid variables that do not depend on the amount of Space, the units of pc are proportional to the ho unit. Since the proportionality constant is unknown, the units of pc will be used. However, if the measurement such as redshift depends on the amount of Space, the units of pc should be used with caution.

Since the hods are discrete, $\epsilon(t)$ is a digital function in time. The hod exerts a tremendous τ to hold the Space in its 2D surroundings. Upon transition, the Space immediately around the hod experiences an instantaneous addition to the amount of Space in the 3D universe. This step function creates a shockwave in the surrounding Space. Space then flows out into the void causing the V of Space to increase and a ρ variation in the V .

If Space has no relaxation time and no maximum gradient, the eruption of Space from a Source would spread out instantly so that ρ would be virtually zero everywhere. Since we observe Space in our region of the universe, the Anthropic Principle implies Space flow must have a relaxation time and a maximum ρ gradient ($\nabla_{\text{m}}\rho$) that is a constant of the universe. Bold face type denotes a vector.

Since Space has a relaxation time, the mirror property in hods is inertia ι per α . A hod is an object with an area α , a surface tension per unit area τ , and inertia per unit area ι . The α , the magnitude of τ , and the ι are the same for all hods and are universal constants.

The τ and ι furnish a resistance to deformation on the hod. Since α is a constant, a deformation of a hod causes the edge to change shape.

In the region around an active Source, $\nabla\rho = \nabla_{\text{m}}\rho$. Space expands into the void where $\rho = 0$ and $\nabla\rho = 0$. Therefore, at some radius r_{sc} from the Source, ρ becomes another function of time and distance from the Source. At r_{sc} , $\nabla\rho$ must be continuous, $\nabla\rho = \nabla_{\text{m}}\rho$ and $\nabla^2\rho > 0$. If $\epsilon(t)$ changes, then r_{sc} changes.

As the Space increases V , Space increases between the hods. The hods move and attain a velocity $d\mathbf{d}_{ij}/dt$ where $\mathbf{d}_{ij}$ is the vector distance from the i^{th} hod to the j^{th} hod. The hod's ι inhibits the movement. Thus, a feedback condition is established wherein the $d\mathbf{d}_{ij}/dt$ is balanced by ι . Therefore, the hods movement is proportional to the total inertia $\iota\alpha$ of the hod and to the velocity of the hod. The momentum $\mathbf{p}_{ij}$ between the of the i^{th} and j^{th} hods is defined as

$$\mathbf{p}_{ij} \equiv K_{\text{p}} \iota\alpha \frac{d\mathbf{d}_{ij}}{dt}, \quad (5)$$

where K_{p} is the proportionality constant and $\equiv$ means "is defined as".

The energy T_{ij} expended by the Space expansion of V upon the hod is

$$T_{ij} = \int \mathbf{p}_{ij} \bullet d\mathbf{d} = K_{\text{p}} \iota\alpha \int \frac{d}{dt} \left| \frac{d\mathbf{d}_{ij}}{dt} \right|^2 dt. \quad (6)$$

Thus,

$$T_{ij} = K_{\text{t}} K_{\text{p}} \iota\alpha \left| \frac{d\mathbf{d}_{ij}}{dt} \right|^2, \quad (7)$$

where K_{t} is the proportionality constant.

The T_{ij} is a scalar quantity and is the kinetic effect of the i^{th} hod on the j^{th} hod. The total T_{ij} energy T_j exerted on the j^{th} hod by all other hods is

$$T_j = \sum_{i \neq j, i=1}^{N_{\text{t}}(t)} T_{ij}. \quad (8)$$

Because the amount of energy added by the movement of $\iota\alpha$ must be the negative of energy due to position and by the inertia which inhibits hod movement, the Principle of Negative Feedback applies. As Space is added

between hods, the energy the hods exert on each other must increase. Since the ρ term of Eq. (2) does not apply to the hod, the loss of energy must be due to $\tau\alpha$. Call this energy between the of the i^{th} and j^{th} hods the potential energy U_{ij} . By the Principle of Negative Feedback, U_{ij} is proportional to $\tau\alpha$ and is a function $f(d_{ij})$ of d_{ij} between hods

$$U_{ij} = -K_{\text{u}} \tau\alpha f(d_{ij}), \quad i \neq j \quad (9)$$

where K_{u} is the proportionality constant and the negative sign is because the hods exert an attraction to each other.

The total U_{ij} energy U_j exerted on the j^{th} hod by all other hods is

$$U_j = \sum_{i \neq j, i=1}^{N_{\text{t}}} U_{ij}. \quad (10)$$

The j^{th} hod's $\iota\alpha$ inhibits the changing $\mathbf{p}_{ij}$. The inertia of the hods results in work W done on the hod in time dt

$$W_j = - \int_{\text{b}}^{\text{e}} \frac{d}{dt} \sum_{i \neq j, i=1}^{N_{\text{t}}} \mathbf{p}_{ij} \bullet d\mathbf{d}, \quad (11)$$

$$W_j = -K_{\text{p}} \iota\alpha \left(d_{\text{e}} \frac{d^2 d_{\text{e}ij}}{dt^2} - d_{\text{b}} \frac{d^2 d_{\text{b}ij}}{dt^2} \right), \quad (12)$$

where subscript "b" and subscript "e" indicate the beginning and ending positions in the time period, respectively.

Therefore, the total energy equation in 3D is Eq. (3) with T_j and U_j is

$$N_{\text{t}} \tau\alpha + \sum_{i \neq j, j=1}^{N_{\text{t}}} (T_j + U_j) - \iiint_{\text{Space}} \rho dV = 0. \quad (13)$$

Implied in Eqs. (3) and (13) is

$$\sum_{i \neq j, j=1}^{N_{\text{t}}} \sum_{i=1}^{N_{\text{t}}} K_{\text{t}} K_{\text{p}} \iota\alpha \left(\frac{d\mathbf{d}_{ij}}{dt} \right)^2 = \sum_{i \neq j, j=1}^{N_{\text{t}}} \sum_{i=1}^{N_{\text{t}}} K_{\text{u}} \tau\alpha f(d_{ij}). \quad (14)$$

"Flatness" is a fundamental condition of the creation of matter and Space. Therefore, the Cartesian coordinate system with four equal angles to define a complete circle is the form of the "flat" measuring system.

Note, $\mathbf{d}_{ij}$ is between hods. The process of summing over all hods provides a reference frame similar to Mach's principle. Also, since there is distance between galaxies, the U_{ij} and T_{ij} of hods in different galaxies includes a heritage from 1D and 2D.

A. Energy continuity equation

Consider an arbitrary volume ΔV bounded by a surface Δs . Space will be in ΔV . Hods, Sources, and Sinks may be in ΔV . Equation (7) implies $T_{ij} = T_{ji}$. Equation (9) implies $U_{ij} = U_{ji}$. Therefore, the total energy E_v in ΔV is

$$\begin{aligned} E_v &= N\tau\alpha + \iiint_{\Delta V} \rho dV \\ &+ \sum_{j=1}^N (T_j + U_j + W_j) \\ &+ \sum_{k=1}^{N_\epsilon} \epsilon_k + \sum_{l=1}^{N_\eta} \eta_l, \end{aligned} \quad (15)$$

where N is the number of hods in ΔV , the integration is over ΔV , N_ϵ is the number of Sources in ΔV , N_η is the number of Sinks in ΔV , ϵ_k is the strength of the k^{th} Source, and η_l is a negative number which is the strength of the l^{th} Sink.

The classical development of an energy continuity equation includes the assumption that energy is neither created nor destroyed. In the CUM, an arbitrary volume may include creation into our universe of energy (Sources) or the destruction from our universe of energy (Sinks).

Following the usual procedure for the development of an energy continuity equation such as the heat equation for the temporal evolution of temperature; remembering hods have no volume and, therefore, the region can be made regular and simply connected by mathematically drawing cuts around the hod discontinuities; and using the Principle of Minimum Potential Energy and the Principle of Feedback and Minimum Action gives the integral form of the energy continuity equation

$$\begin{aligned} \iiint C \frac{d}{dt} E_v dV &= S \iint \nabla \rho \bullet \mathbf{n} ds \\ &+ S_u \iint \nabla \left| \sum_{i \neq j, j=1}^N U_j \right| \bullet \mathbf{n} ds \\ &+ S_t \iint \nabla \left| \sum_{i \neq j, j=1}^N T_j \right| \bullet \mathbf{n} ds \\ &+ S_w \iint \nabla \left| \sum_{i \neq j, j=1}^N W_j \right| \bullet \mathbf{n} ds \\ &+ \iint [\tau \alpha N_{\text{tuw}}(t) \mathbf{v}_n] \bullet \mathbf{n} ds \\ &+ \iiint \left(\sum_{k=1}^{N_\epsilon} \epsilon_k + \sum_{l=1}^{N_\eta} \eta_l \right) dV, \end{aligned} \quad (16)$$

where: (1) The S is a constant of proportionality. Call S the conductivity. S is the ability of energy to flow

and is assumed to be independent of position or direction. S is a factor that slows the flow of Space. The idea of their being some kind of viscosity or friction associated with Space would require the expenditure of energy. Since Equation (13) lacks such a term, Space energy dissipates without loss of total energy. Therefore, there is a relaxation time associated with Space. This means that the stress created by the sudden entry of Space upon a transition must relax asymptotically with time. The compression of the transition is dissipated by the flow of Space into the void. (2) The S_u is a constant of proportionality. Call S_u the potential conductivity. S_u is the ability of energy to flow and is assumed to be independent of position or direction. (3) The S_t is a constant of proportionality. Call S_t the kinetic conductivity. S_t is the ability of kinetic energy to flow and is assumed to be independent of position or direction. (4) The S_w is a constant of proportionality. Call S_w the work conductivity. S_w is the ability of work energy to flow and is assumed to be independent of position or direction. (5) The $N_{\text{tuw}} = N + \sum_{j=1}^N (T_j + U_j + W_j) |_{T_j, U_j, W_j}$ is the number of hods which leave the ΔV with the energy associated with each hod and $|_{T_j, U_j, W_j}$ means that T_j , U_j , and W_j are constant with respect to the differentiation. (6) The $\mathbf{v}_n$ is the velocity vector of the N_{tuw} .

Applying Gauss's divergence theorem to the surface integrals of Equation (16), and noting that the volume could be chosen to be a region throughout which the integrand has a constant sign yields the differential form of the energy continuity equation

$$\begin{aligned} \frac{dE_v}{dt} &= D^2 \nabla^2 \rho + \frac{S_u}{C} \nabla^2 \left| \sum_{i \neq j, j=1}^N U_j \right| \\ &+ \frac{S_t}{C} \nabla^2 \left| \sum_{i \neq j, j=1}^N T_j \right| + \frac{S_w}{C} \nabla^2 \left| \sum_{i \neq j, j=1}^N W_j \right| \\ &+ \nabla \bullet [\tau \alpha N_{\text{tuw}}(t) \mathbf{v}_n] \\ &+ C^{-1} \left(\sum_{k=1}^{N_\epsilon} \epsilon_k + \sum_{l=1}^{N_\eta} \eta_l \right), \end{aligned} \quad (17)$$

where $D^2 = S/C$ and is called the diffusivity constant of Space.

The presence of Sources and Sinks leads to a non-homogeneous equation. Note the similarity of this equation to the diffusion equation for the temporal evolution of the concentration of chemical species in a fluid and the heat equation for the temporal evolution of distributions of temperature. Like the molecular diffusivity and thermal diffusivity, D^2 has its origins in the dynamics and interactions that transfer energy.

B. Forces

The ρ terms and U_j term in Eq. (15) change by spatial movement and are called impelling forces since they cause

an energy change. The $N\tau\alpha$ term, the W_j term, and T_j term in Eq. (15) change by temporal differences and are called nurturing forces since their movement carries energy.

1. Space

Define a volume V containing Space for the distances being considered and throughout which the terms of Eq. (17) except ρ terms are constant. The force of Space at a point $\mathbf{F}_{\text{sp}}$ is defined as

$$\mathbf{F}_{\text{sp}} \equiv -D^2 \nabla \rho, \quad (18)$$

where the negative sign means the force is directed opposite to increasing ρ .

Because there is no shear stress in Space, the force exerted by Space is exerted only normal to surfaces. Consider a cylinder of cross section ds around a regular, simply connected volume with the ends of the cylinder normal to $\mathbf{F}_{\text{sp}}$. The ds has a difference of force $\Delta \mathbf{F}_s$ on each end of the cylinder. Allow the height of the cylinder to become arbitrarily small. The $\Delta \mathbf{F}_s \rightarrow d\mathbf{F}_s$ and

$$d\mathbf{F}_s = -D^2 (\mathbf{n} \bullet \nabla \rho) ds, \quad (19)$$

where $\mathbf{n}$ is the outward unit normal of ds .

Integrating Eq. (19) over a surface, the Space force on the surface $\mathbf{F}_s$ becomes

$$\mathbf{F}_s = -D^2 \iint \nabla \rho \bullet \mathbf{n} ds. \quad (20)$$

The force of Space on a surface is proportional to the cross section of the surface perpendicular to $\nabla \rho$.

The ρ due to Sources and Sinks can be calculated from Eq. (17). If all terms of Eq. (17) are constant except the ρ , ϵ , and η terms, the average η of each Sink remains constant over a long period of time, and the average ϵ of each Source, each galaxy, remains constant over a long period of time, then Eq. (17) can be solved for a steady-state condition. The range of distance from Sources and Sinks will be assumed to be sufficient such that the transition shock is dissipated and $r > r_{\text{sc}}$. We can consider ρ at a point to be the superposition of the Space effects of each Source and Sink. Therefore, for ρ from the j^{th} galaxy and k^{th} Sink at the i^{th} point (ρ_i), Eq. (17) becomes

$$\frac{d}{dt}(r_{ij}\rho_i) = D^2 \nabla^2 (r_{ij}\rho_i), \quad (21a)$$

$$\frac{d}{dt}(r_{ik}\rho_i) = D^2 \nabla^2 (r_{ik}\rho_i) \quad (21b)$$

in spherical coordinates, where r_{ij} is the distance from the j^{th} galaxy Source and r_{ik} is the distance from the k^{th} Sink to the i^{th} point, and ρ_i depends entirely on r_{ij} and r_{ik} .

The boundary conditions are:

$$\rho_i(r_{ij}, 0) = \text{a function of distance, only}, \quad (22a)$$

$$\rho_i(r_{ij}, t) \rightarrow 0, \text{ as } r \rightarrow \infty, \quad (22b)$$

$$\rho_i(r_{ik}, 0) = \text{a function of distance, only}, \quad (22c)$$

$$\rho_i(r_{ik}, t) \rightarrow 0, \text{ as } r \rightarrow \infty. \quad (22d)$$

This is analogous to the problem of heat flow in a spherical solid with a continuous Source of strength ϵ_j at $r_{ij} = 0$ or with a continuous Sink of strength η_k at $r_{kk} = 0$. By superposition the total $\rho_i(x, y, z)$ may be calculated [50] for r_{ij} and r_{ik} outside the $\nabla \mathbf{m}\rho$ volume of the Sources and Sinks as

$$\rho_i(r) = \frac{1}{4\pi D^2} \left(\sum_{j=1}^{\text{all galaxies}} \frac{\epsilon_j}{r_{ij}} - \sum_{k=1}^{\text{all Sinks}} \frac{|\eta_k|}{r_{ik}} \right). \quad (23)$$

At $r_{ij} = r_{\text{sc}}$ of the j^{th} galaxy, the effect of the other Sources and Sinks is very small. Therefore,

$$\nabla \rho_i = \frac{\epsilon_j}{4\pi D^2} \nabla r_{\text{sc}j}^{-1} = \nabla \mathbf{m}\rho, \quad (24)$$

where $r_{\text{sc}j}$ is the r_{sc} for the j^{th} galaxy.

At the j^{th} Source, the nearly instantaneous transition of the hods and Space into 3D creates a shock wave zone of radius $r_{\text{sw}j}$, where the transition forced $\nabla \rho$ is greater than $\nabla \mathbf{m}\rho$. The assumption of constant D is no longer valid. In the region $r_{\text{sw}j} < r_{ij} < r_{\text{sc}j}$, ϵ_j can be considered to be a time average of the nearly digital transitions and $\nabla \rho_i = \nabla \mathbf{m}\rho$.

At the Source of the j^{th} galaxy $r_{ij} = 0$, $\rho_i(r_{ij} = 0)$ is a maximum value $\rho_{\text{max}j}$ and $\nabla \rho_i$ and $\nabla^2 \rho_i$ are discontinuous. For periods of the time required for Space to flow to $r_{\text{sc}j}$, ϵ_j can be considered a constant. All the Space from the Source is flowing through the surface of the sphere of radius $r_{\text{sc}j}$. Since the volume of a sphere $\propto r^3$ and ϵ is a volume per unit time, $\epsilon \propto r$. Further, this proportionality holds for any radius determined by ϵ . If ϵ_j increases, $r_{\text{sc}j}$ increases proportionally as suggested by Eq. (24). Therefore,

$$r_{\text{sc}j} = \frac{K_{\text{sc}}}{|\nabla \mathbf{m}\rho|} \epsilon_j, \quad (25)$$

where K_{sc} is the proportionality constant which depends on how D varies in the $\nabla \mathbf{m}\rho$ zone around the Source and $\rho_{\text{max}j} = K_{\text{sc}} \epsilon_j$.

The force $\mathbf{F}_{\text{ts}}$ exerted on an assembly of hods by the Space force due to Sources and Sinks is calculated by combining Eq. (20) and (23) and integrating over the cross section of the assembly yields

$$\mathbf{F}_{\text{ts}i} = G_s m_s \left(\sum_{j=1}^{\text{all galaxies}} \frac{\epsilon_j}{r_{ij}^3} \mathbf{r}_{ij} - \sum_{k=1}^{\text{all Sinks}} \frac{|\eta_k|}{r_{ik}^3} \mathbf{r}_{ik} \right), \quad (26)$$

where m_s is the Space effective cross section of the assembly of hods and $G_s = (4\pi)^{-1}$.

2. Interaction of Space and hods

The hod in 2D is transformed into 3D because of the energy pressure of surrounding hods. The transition releases this energy into 3D. The τ acts on Space uniformly across the surface of the hod and exerts an attractive pressure on Space normal to the surface of the hod. Since the hods in 2D are closed loops in a plane, the hods arrive in 3D as surfaces oriented flat-to-flat. At the transition, $V_I = 2d_I\alpha$ where d_I is the initial distance that Space extends from a hod's surface. At the radius r_h of the hod from the Source, Space has expanded and the $\rho_i(r_h) < \rho_I$. However, within $0 < r_h < r_{sw}$, $\nabla\rho = \nabla_{\mathbf{m}}\rho$ and $\rho_h < \rho_i$ where ρ_h is the maximum Space density the hod can attract to its surface. All the Space energy in the shock wave zone is within a zone with $\nabla\rho = \nabla_{\mathbf{m}}\rho$. Therefore, all the $\rho_I V_I$ is "bound" to the hods $\rho_I V_I = \tau\alpha = \tau_b\alpha$, where τ_b is the amount of surface tension used to bind Space to the hod.

Define a Cartesian coordinate system in a volume V with the origin at the center of a circular hod with diameter a , with the z -axis perpendicular to the plane of the hod, and with the x -axis and y -axis in the plane of the hod.

At a radius r_h from the Source outside the shock wave zone, the maximum distance d_h (ho) from the hod surface along the z axis within which $\nabla\rho = \nabla_{\mathbf{m}}\rho$ is directed away from the hod surface has a value

$$d_h = \frac{\rho_h - \rho_d}{|\nabla_{\mathbf{m}}\rho|}, \quad \rho_d = \rho_i(d_h) < \rho_h, \quad r_{sw} < r_h, \quad (27)$$

where ρ_d is the Space density at d_h .

Hods that are within $2d_h$ of each other form an assembly of hods within a volume with a $\rho = \rho_d$ equipotential surface. For $r_{sw} < r_h \leq r_{sc}$, $\nabla\rho = \nabla_{\mathbf{m}}\rho$ everywhere.

For $r_h > r_{sc}$, volumes more than d_h from a hod surface have $\nabla\rho < \nabla_{\mathbf{m}}\rho$, $\rho = \rho_i$, Eq. (27) applies, and $\tau_b\alpha < \tau\alpha$. Therefore, there is an amount of "free" surface tension energy τ_f such that

$$\tau\alpha = \tau_b\alpha + \tau_f\alpha \quad (28a)$$

$$\tau_f = d_h(\rho_h - \rho_d), \quad r_h > r_{sc}, \quad (28b)$$

$$\tau_b = 2d_I - \tau_f, \quad (28c)$$

$$\tau_f = 0, \quad r_h \leq r_{sc}. \quad (28d)$$

The τ_f is exerted on ρ_i . For the remainder of this Paper, unless otherwise stated, the region under consideration is with $\rho_i < \rho_h$. Those hods that are oriented flat-to-flat emerge from the r_{sc} zone as assemblies of hods. Other hods become independent.

Since the energy at the Source upon transition is high and since we observe Space to be more extensive than the hods, $0 < d_h \ll 1$ ho.

Consider the method of conjugate functions of a complex variable with a transformation

$$t = \frac{a}{2} \cosh(w), \quad (29)$$

where $t = x + iz$ and $w = u + iv$. Therefore,

$$\frac{x^2}{\cosh^2 u} + \frac{z^2}{\sinh^2 u} = \frac{a^2}{4}, \quad (30)$$

$$\frac{x^2}{\cos^2 v} - \frac{z^2}{\sin^2 v} = \frac{a^2}{4}. \quad (31)$$

Geometrically, the strip in the w -plane between the lines $v = 0$ and $v = \pi$ transforms into the entire t plane. The line at $u = 0$ between $v = 0$ and $v = \pi$ transforms to the surface of the hod. The lines $v = \text{constant}$ are transformed into confocal hyperbolae lines which are streamlines of the ρ_{ij} field where ρ_{ij} is the ρ at the j^{th} point due to the i^{th} hod. The lines $u = \text{constant}$ are transformed into confocal ellipses with the foci at the ends of the hod. Since τ acts normally to the surface of the hod, $u = \text{constant}$ are equipotential lines of ρ_{ij} caused by τ [see Eq. (9)].

If the hod is circular around the z -axis, the ρ_{ij} streamlines and equipotential lines are also circular around the z -axis. The ρ_{ij} equipotential lines form oblate spheroids with the minor axis along the z -axis.

Consider a point at a large distance from the hod, $r \gg a$, the u of Eq. (30) is very large. Equation (30) becomes $x^2 + z^2 = d^2$ where d is the distance from the center of the hod. At large d the ρ_{ij} equipotential lines are concentric spheres. The ρ_{ij} streamlines are the radii of the spheres.

Since the equipotential volume surrounded by ρ_d is an oblate spheroid, the distance d_x (in ho) in the plane of the hod to $\rho = \rho_d$ is

$$d_x = \sqrt{d_h^2 + 0.25} - 0.5, \quad (32)$$

which is much smaller than a .

Therefore, the $\rho = \rho_d$ equipotential may be approximated as a cylinder with height $2d_h$ and end area $= \alpha$.

If $\nabla\rho_s > 0$ across a hod, the $\nabla\rho$ on one side of the hod at an infinitesimal distance dd from the $\rho = \rho_d$ surface $\nabla\rho_{s1}(d + dr)$ differs from the other side $\nabla\rho_{s2}(d - dr)$. Since the $\nabla\rho$ is limited to $\nabla_{\mathbf{m}}\rho$, the force of Space from Eq. (19) is transferred to the hod surface. Therefore, the net force $d\mathbf{F}_s$ of Space on an element of surface ds of the hod is

$$d\mathbf{F}_s = D^2(\mathbf{n} \cdot \nabla\rho_{s2})ds - D^2(\mathbf{n} \cdot \nabla\rho_{s1})ds, \quad (33a)$$

$$d\mathbf{F}_s \approx -D^2(\mathbf{n} \cdot \nabla\rho_s)ds, \quad (33b)$$

where $\nabla\rho_s$ is considered approximately constant over the small $2d$ distance. This approximation fails in volumes close to r_{sc} .

Unless otherwise stated, when distances greater than one ho are implied, $d \ll 1$ ho and $\rho = \rho_s$ will be considered to be at the hod surface.

Since the hod is supple, the application of $d\mathbf{F}_s$ on ds of the hod in the x - y plane will cause ds to move normal to the hod surface. If $d\mathbf{F}_s = 0$ everywhere else, a deformation, du , of the hod surface normal to hod surface will occur. Since α and τ are constant, the edge of

the hod must change shape. A small deformation allows movement in only the z direction. The edge stays in the x - y plane. However, since edges of a free, individual hod are not constrained and τ is very strong, the hod membrane does not oscillate normal to the hod surface as it would for a stretched membrane. Therefore, $d\mathbf{F}_s$ will cause the hod to tend to conform to the $\nabla\rho_s$ streamlines. Since Space has a relaxation time, the movement of hods will cause the streamlines to be continually shifting. A slight delay, caused by ι and by S , in adjusting to a change in the streamline will allow interesting effects. If $\nabla\rho_s \bullet \mathbf{n} \neq 0$ for a brief instant on only one edge of the hod and if the other edge experiences a force parallel to the surface $\nabla\rho_s \bullet \mathbf{n} = 0$. The hod will travel along the streamline at the maximum allowed velocity. If the streamline the hod is following has a very sharp bend, such as at the edge of another hod where the hod is much stronger than anywhere except near the Source, the inertia energy of the hod will let it continue. For an instant, $\nabla\rho_h \bullet \mathbf{n} \neq 0$. The hod will slow or even reverse direction. This allows the transfer of energy between hods.

If the $\nabla\rho$ field of Space varies little over the surface of the hod, the deformations become a general shaping of the hod causing the hod to tend to conform to the force lines, or streamlines, of ρ . The hod will move to make $\nabla\rho \bullet \mathbf{n} = 0$ on the surface of the hod.

If $\nabla\rho \bullet \mathbf{n} = 0$ from some direction, the hod will experience zero Space impelling force from that direction. If the $\nabla\rho$ field changes little over a range of several hod diameters, small deformations with $\nabla\rho \bullet \mathbf{n} \neq 0$ will cause the hod to have a forward inertia.

3. Gravity

Call $\mathbf{F}_{gk}$ the force of gravity on the k^{th} hod,

$$\mathbf{F}_{gk} \equiv -\frac{S_u}{C} \sum_{k \neq j=1}^{N_t} \nabla U_{kj}, \quad (34)$$

where the negative sign means the force is directed opposite to increasing U and the sum is over the other (j^{th}) hods.

Define a volume V containing Space for the distances being considered and throughout which the terms of Eq. (17) except the U terms are constant. By the Principle of Superposition, this allows the examination of just one force. Equation (17) can be solved for a steady-state condition. We can consider ρ at a point to be the superposition of the effects of each hod. Therefore, Eq. (17) at the i^{th} point due to the j^{th} hod becomes

$$\frac{d}{dt} (d_{ij} U_j) = \frac{S_u}{C} \nabla^2 (d_{ij} U_j) \quad (35)$$

in spherical coordinates.

The boundary conditions are:

$$U_i(d_{ij}, 0) = \text{a function of distance, only}, \quad (36a)$$

$$U_i(d_{ij}, t) \rightarrow 0 \text{ as } r \rightarrow \infty. \quad (36b)$$

Analogous to Section II B 1,

$$U_j = -\frac{K_u C \tau_f \alpha}{4\pi S_u d_{ij}}. \quad (37)$$

The $|U_{ij}|$ is highest in a volume with a given $\tau_f > 0$ value when the k^{th} and j^{th} hods have a shared $\rho = \rho_d$ oblate spheroid surface and are orientated flat-to-flat. Since d_{ij} is the distance to the center of the hod, $d_{ij} \neq 0$ and a singularity problem is non-existent.

Chose two assemblies of hods where N_t is the number of hods in one assembly and N_s is the number of hods in the other assembly. Let each assembly have a very small diameter compared to the distance $\mathbf{r}_{ts} = \langle \sum_{j=1}^{N_t} \sum_{k=1}^{N_s} d_{kj} \rangle$ where $\langle \rangle$ means ‘‘average of’’. The force of gravity of the hods in an assembly is greater on each other than the force of gravity of hods in the other assembly. Therefore, the hods of an assembly will act as a solid body relative to the other assembly of hods. Because hods act on Space, Space acts on Space and Space acts on hods, only the local τ_f for each hod determines the $N \tau_f \alpha$ on the hod. Therefore, the force of gravity $\mathbf{F}_g$ on each assembly due to the other assembly is

$$\mathbf{F}_g = -\frac{(K_u \alpha)^2}{4\pi r_{ts}^3} (N_s \tau_{fs}) (N_t \tau_{ft}) \mathbf{r}_{ts}, \quad (38)$$

where τ_{fs} is the τ_f for the assembly with N_s hods and τ_{ft} is the τ_f for the assembly with N_t hods.

For the simplified case of two symmetric distribution of hods with a distance $\mathbf{r}_{12}$ between them and with $\tau_{fs} = \tau_{ft}$

$$\mathbf{F}_g = -G \frac{m_{g1} m_{g2}}{r_{12}^3} \mathbf{r}_{12}, \quad (39)$$

where $G = (4\pi)^{-1} (K_u \alpha)^2$, $m_{g1} = N_t \tau_{ft}$, and $m_{g2} = N_s \tau_{fs}$. The experimental determination of m_{g1} and m_{g2} must use a method depending on U_{ij} and no other factor such as motion which will involve $\iota \alpha$ forces.

Note, in volumes within a sphere with a radius of r_{sc} from a galaxy’s Source, $\tau_f = 0$ and, therefore, $\mathbf{F}_g = 0$ and $m_g = 0$. In volumes with $r_{sc} < r_h$, τ_f increases as r_h increases.

C. Particles

Define a particle to be an assembly of hods surrounded by a $\rho = \rho_d$ equipotential surface (ES) that forms a connected volume around the hods in the particle within which Space is totally bound to the hods by τ_b . Therefore, ρ_s outside the ES equals ρ_d within the ES of the particle. If the ES is simply connected, the particle is

called a simple particle and the distance for gravity potential calculations is to the center of the particle. The potential on a surface element of the ES is determined by the distribution of hods inside the surface. However, the volume occupied by the bound Space is the same for each hod. Therefore, the hods are uniformly distributed over the volume inside a simply connected ES. Since the hods of a simple particle have a minimum distance between them, the hods are tightly bound and respond as a single entity. By the Principle of Minimum Potential Energy, the hods of a simple particle have a more stable structure with a maximum $N\tau_b\alpha$ per unit surface area ratio (P/S).

Hods are unable to join outside the Source region to form a particle but may be able to join an existing particle. The existing particle must have sufficient $N(\tau_f\alpha)$ to overcome the $\iota\alpha$ of the joining hod. Also, this implies there is a minimum number $N_{\min}$ of hods which must be combined before other hods may join it outside the Source region.

When the distance between the flat of the hods $\leq 2d_h$, the $\nabla\rho$ between them will change sign and become $\nabla_m\rho$ which will cause a net repulsive force between hods. Therefore, the distance between the flat of the hods will stabilize at a distance $2d_h$ when the repulsion due to $\nabla_m\rho$ will equal F_g . If the distance between the flat of the hods $> 2d_h$, the hods will be independent.

Also, since $d_h \ll 1$ ho, hods may form in a column arrangement. Call this type of structure Family 1 (F1) Type 1 (T1) structures with the length $L = 2N_c d_h$ where N_c is the number of hods in the F1T1 structure. The potential can become large enough that hods can be added without being in a Source r_{sc} zone. The P/S of the F1T1 structure will be least when $2N_{\min}d_h = 1$ ho. The addition of hods to the column also produces stable, cylinder structures since each additional hod is oriented flat-to-flat. Therefore, the addition of hods to column length decreases the P/S. However, for a large number of hods, the outer hods are easier to remove. The direction parallel to the surface has $\nabla\rho \bullet \mathbf{n} = 0$ so $F_s = 0$ and maximal speeds can be achieved in a direction parallel to the surface. This structure is a photon. The direction along the axis has a one-hod surface. Hence, F_s along the cylinder axis is as if only one hod were present.

If the inertia and external forces cause the hods in the F1T1 to start to lose overlap, the sharp $\nabla\rho$ field at the edges of hods will push them back. The total overlap condition is a minimum potential energy position.

Any deformation in the surface from a ripple in Space will tend to bend the hod surfaces to lose total overlap. Since this causes a force to reestablish overlap, this structure is much more rigid than a single hod. Because this structure is rigid, a $\nabla\rho$ across the structure can impart a torque on the particle.

As more hods are added, the $N_c(\tau\alpha)$ increases. The energy of the photon $h\nu$ is

$$h\nu = N_c(\tau\alpha). \quad (40)$$

Ripples in Space will cause ripples in the surface of the outer hod. This will slow the photon structure and cause a shearing F_s on the outer hod. Thus, if $N_c > N_{\min}$ in a photon, the photon can lose hods or gain hods.

Two cylindrical photons may join as side-by-side cylinders caused by F_g . Parallel to the surface, $\nabla\rho_s \bullet \mathbf{n} = 0$ for both.

Also, two cylindrical photons may join in a T formation if $L > 1$ ho as the Principle of Change requires. By extension, if $L \approx 2$ ho, there may be other cylinders added to form II, III, etc. structures. This requires the photons in the vertical columns to have approximately, if not exactly, the same number of hods. Otherwise, one or more of the photons will experience a $\nabla\rho \bullet \mathbf{n} \neq 0$. There will still be a direction where $\nabla\rho \bullet \mathbf{n} = 0$. Call this structure a F1, Type 2 (T2) structure. Perhaps this is the structure of a lepton.

Consider a F1T2 structure made of several columns and a cross member at each end. If an end column does not overlap the cross member column, an extremely unstable condition exists. The end column would be freed. If a cross member is longer than the width of the row of columns, the particle could acquire another column. The most stable structure would be when L is an integer multiplier I of the average diameter of a ho

$$L = 2I\left(\frac{\alpha}{\pi}\right)^{1/2}. \quad (41)$$

Similarly, a third photon can join the T structure so that the three are mutually orthogonal, as the Principle of Repetition allows. However, $\nabla\rho \bullet \mathbf{n} \neq 0$ in all directions. Call this structure a F1, Type 3 (T3) structure. A cross cylinder on a cross cylinder of a F1T2 structure makes the F1T3 box structure very dense and very stable since there is little “free” surface (unattached hod faces) compared to the $\tau_f\alpha$ energy contained and P/S is minimal. Perhaps this is the structure of quarks, quark stars, and black holes.

In small structures, the F1T3 cube has the least P/S. However, this allows “corners”. In larger structures, a rounding effect such as found in crystal structures produces a more spherical ES which will have a lower P/S. This could be a mechanism to fracture some combinations (particles) and cause faceting in all combinations like in crystals.

Another possible structure of hods with a position of least P/S and stable $\nabla\rho$ field is with each inserted through and normal to each other (see Fig. 2). Call the particles formed with this structure Family 2 (F2) particles. The particle so formed has a direction along the line where they join where $\nabla\rho \bullet \mathbf{n} = 0$ can occur. These may be the Muon family of particles.

By a similar argument, with less probability, three hods may come to form a particle with three mutually orthogonal hods (see Fig. 3). Call the particles formed with this structure Family 3 (F3) particles. These may be the Tau family of particles.

However, a 4th mutually orthogonal hod construction in 3D outside the immediate area of a Source is not fea-




FIG. 2: Diagram showing the structure of the Family 2 (Muon family) fundamental particles.




FIG. 3: Diagram showing the structure of the Family 3 (Tau family) fundamental particles.

sible since one of the hod's surfaces will face another hod surface.

Therefore, because the hod defines 3D, there are three, and only three, families of particles.

In our volume of the universe, particles of the same type have consistent mass. Particles become larger from gravitational attraction of additional hods as the Principle of Repetition allows. The observed consistent mass of like particles implies the existence of the Principle of Negative Feedback to limit the size. For T2 and T3 particles, if $L > I(\alpha/\pi)^{1/2}$ from Eq. (41), attracted hods may be added to the particle. As noted previously, the attracted hod will be traveling at the maximum speed which implies a maximum T . Thus, the attracted hod has energy

great enough to break the gravitational bond and facet the outlying hods from the particle. If $L < I(\alpha/\pi)^{1/2}$, the energy of the attracted hod may cause the column to facet.

In a volume with slow changing ρ_i , $\rho_d = \rho_i$. Therefore, L is a function of ρ_i . In a volume with rapidly changing ρ_i , ρ_d may lag the change in ρ_i because τ_b may be absorbing or releasing Space. Such may be the situation for neutron stars, possible quark stars, and black holes, which exist near the Source of the Galaxy.

If the particles are to have more hods (energy/mass) as the Anthropic Principle requires, there must be another binding mechanism less sensitive to faceting as the Principle of Change requires. The flat side of the end hods on a photon would be attracted to and could be bound to other particles. If quarks are the largest Family "type" particles, baryons must be quarks and leptons joined by another means such as directly to each other or by photons. The particle will be in a connected volume, but not a simply connected volume. By the Principle of Repetition, the atom is electrons, neutrons, and protons joined by photons. The photon is similar to a rod in these structures.

D. Sink characteristics

If a Source ceases to erupt or if some matter is repelled to the low $\nabla\rho$ distances between Sources, gravity will cause the hods to coalesce. In a volume where $\nabla\rho$ becomes near zero, the attraction of hods will cause a high hod density. If the number of hods is greater than in a black hole in this intergalactic volume, the intense force on the supple hods may cause the hods to form spheres like bubbles. With increased pressure from a huge number of hods, the hod can go only out of our universe and turn into a four dimensional 4D object like 2D turned to 3D.

Since the Sink requires mass to form, the Sink's age is considerably less than galaxies. Therefore, the delay before matter and Space is sent into the 4D allows 3D to accumulate mass.

The Sink requires a minimum mass to send hod bubbles into 4D. Therefore, the mass M_{sink} around the Sink has an upper limit. If the Sink is sending hods to 4D, then a negative feedback condition is formed and the rate η of transition of the hods is dependant on the rate of accretion of hods. Therefore, $\eta \propto M_{\text{sink}}$.

The $\nabla\rho$ acts in the same direction as the gravity force. The Sink can send hods to 4D if the hods are close. Therefore, the $\nabla\rho = \nabla_m\rho$ around the Sink like around the Source.




FIG. 4: Diagram depicting the regions of a galaxy in relation to a hypothetical rotation curve. The lines in the Saturation Region depict the different shapes possible.

III. GALAXY MODEL

A. General

For the purpose herein, a galaxy's RC is divided into five regions [29] as the r from the Source increases (diagrammed in Fig. 4), a center region (CR) [31, 32, 41], a Keplerian region (KR) [15, 16], a rising region (RR) which is in the outer bulge region, a saturation region (SR) which is in the disk region, and a detached region (DR) which is outside the galaxy (herein, collectively referred to as the "lettered regions"). The distinct nature of each region of the RC implies different physics exists among the regions. Therefore, some of the parameters of the equation of motion have suddenly become more or less significant. Each of the lettered regions is separated by a transition region in which the physics is changing from one region to the next. The CR and KR are separated by a transition region T_{c-k} [10, 15, 21, 24]. The KR and RR are separated by a transition region T_{k-r} in which the RC has a slight decline followed by a rapid rise. A similar transition region T_{r-s} appears between the RR and SR. Particles in DR are not in orbit around the galaxy center.

Dividing into regions with different, predominant physics has been done for models of other phenomena. For instance, the physics of semiconductors uses such a model to describe differing current-voltage characteristics of the differing regions of transistor operation [51].

The $\tau_f = 0$ out to r_{sw} and only unbound hods and photons exist near the Source. By Eq. (25) the size of ϵ is proportional to r_{sc} and, therefore, to the diameter of the Source region. As the hods and photons travel outward beyond r_{sc} , $\tau_f > 0$ and particles form.

As the particles' radius r from the Source increases, larger particles can form. Outward from the Source, photons coalesce to form larger particles. In the Big Bang model, the physics of the formation of particles is outward in time. In the CUM, the physics of formation of subatomic particles is outward in distance from the Source. Eventually, hydrogen forms and is pushed outward by the Space force. As hydrogen coalesces, larger atoms are formed which fall back to the center as the gravitational force becomes larger than the Space force. Gravity attracts the densest particles into the center. As the particle approaches r_{sc} , the declining τ_f and L causes the T2 and T3 particles to disintegrate to photons and re-radiate outward.

Consider a ΔV containing a test particle with N hods such as a star or hydrogen gas in a galaxy. The inertial mass m_i is $N\alpha$, the gravitational mass m_g is $N\tau_f\alpha$, and the cross section mass m_s is the effective cross section subject to the Space force of the hods in ΔV . For simplicity, the number of hods in the particle is assumed to be constant over time. Since stars are changing the number of hods by radiation emission and are changing the m_s/m_i ratio through changing elemental composition, this assumption is most nearly true when the ΔV contains hydrogen gas. Therefore, the Hf RC is preferred, where possible, to reflect the forces influencing v .

Assume the particle in ΔV originated in the galaxy ($d_b = 0$) and the particle in ΔV is in equilibrium ($E_v = N\tau_f\alpha$). The coordinate system center is placed at the Source and is aligned in a constant orientation relative to our view. The velocity and rotation of the galaxy relative to the universe is not part of v . Combining Eqs. (23), (37), (8), and (12) into Eq. (15), rearranging terms, and considering only radial terms yields

$$v^2 = r\ddot{r} + \frac{m_g}{m_i} \frac{GM}{r} - \frac{m_s}{m_i} \frac{G_s\epsilon}{r} - \mathbf{r} \bullet \mathbf{a}_o, \quad (42)$$

where the double dots above the r mean the second derivative with respect to time, $\ddot{r}$ is the radial acceleration of the test particle, G is the gravitational constant, G_s is the cross section constant, M is the mass of the galaxy inside a sphere with a center at the Source and a radius of r , ϵ is the Source strength of the galaxy,

$$\mathbf{a}_{oi} = \sum_{j=1}^{\text{other galaxies}} \frac{m_g GM_j - m_s G_s \epsilon_j}{m_i r_{ij}^3} \mathbf{r}_{ij} + \sum_{k=1}^{\text{Sinks}} \frac{m_g GM_k + m_s G_s |\eta_k|}{m_i r_{ik}^3} \mathbf{r}_{ik}, \quad (43a)$$

$$\mathbf{a}_o = \sum_{i=1}^N \mathbf{a}_{oi}, \quad (43b)$$

M_j is the total mass of the j^{th} galaxy, ϵ_j is the Source strength of the j^{th} galaxy, $\mathbf{r}_{ij}$ is the distance from the i^{th} test particle to the j^{th} galaxy's Source, M_k is the total mass of the k^{th} Sink, η_k is the Sink strength of the k^{th}

Sink, m_i is the inertial mass of the test particle, and $\mathbf{r}_{ik}$ is the distance from the i^{th} test particle to the k^{th} Sink.

If $\mathbf{a}_o \neq 0$ in a galaxy, then the center of mass may be slightly displaced from the Source. The v is measured along the major axis from our view. Therefore, the $\mathbf{r}$ is directed along the major axis. The form of Eq. (43) for baryons is of a Newtonian force with an effective mass less than the total mass in a galaxy. Also, the force on F1T1 (photons) particles passing by the galaxy differs because of the orientation of the particle relative to the galaxy.

In our part of the Galaxy, $m_i = m_g$ to within one part in 10^{11} [52]. However, the measurement of m_g depends on $\mathbf{F}_g$ and, therefore, τ_f that depends on ρ . The ratio of the diameter of the earth divided by the distance from the center of the Galaxy suggests much greater precision is required for earth bound measurements. Therefore, the m_g/m_i ratio will be explicitly stated in this Paper where applicable.

The notation convention used herein is as follows. The italic, normal size letter will denote the parameter, as usual. The upper case M means the total mass inside the radius denoted by the sub letters. The sub letters denote the radius range by region identification. The use of “max” or “min” in the sub letters will denote the maximum value or minimum value the parameter has in the region, respectively. For example, the radius r_k (kpc) in the KR which is between the radius r_{tckmax} at the maximum extent of the T_{c-k} region and the radius r_{kmax} at the maximum extent of the KR region. Hence, $r_{\text{tckmax}} < r_k \leq r_{\text{kmax}}$. The notation M_{kmax} means the total mass inside the sphere with radius r_{kmax} . This is the total mass of the CR, T_{c-k} , and KR regions.

The CUM suggests the change in physics among the lettered regions derive from a change in the particle m_s/m_i , in the forces ($\mathbf{F}_{ts}$ and $\mathbf{F}_{tg}$), and in $\mathbf{r} \bullet \mathbf{a}_o$. The lettered regions change is shared by all galaxies and, hence, may be used to derive relations among galaxies.

The following sections discuss each region. Some sections have data that can be compared to the equations developed.

B. CR to KR

The CUM posits that the radius of the CR is less than a few light hours around a Source at the center of a galaxy and is less than r_{sc} . At a radius of less than r_{sw} , there is no v and the problem of a singularity in Eq. (42) at $r = 0$ is nonexistent. Since τ_f decreases as the r decreases below r_{sc} , the gravitational term is decreasing and the ϵ term dominates to repel particles.

The unbound hods and photons in the CR are repelled away from the Source. The ϵ/r close to the Source is large. Therefore, only exceptionally dense particles with very small m_s/m_i are closest to the Source. Black holes which may be F1T3 particles and other very dense matter fall into the T_{c-k} of galaxies. All other matter is repelled

outward. Black holes form a shell around the CR [29, 41].

Black holes may increase m_s/m_i by collision and merger or may lose angular momentum such that they fall into the $r < r_{sc}$ zone. The result of the $L \rightarrow 0$ and $\tau_f \rightarrow 0$ is a multitude of X-ray point sources without accompanying large variation at longer wavelengths near the center of the Milky Way. Note the force of gravity must be reduced to allow the photons to separate in a burst. The diffuse component of the X-ray sources may be other low m_s/m_i particles such as iron nuclei or neutron stars with F1T3 centers [20, and references therein]. At and near r_{sc} the brightness of the radiation will increase relative to star luminosity due to the reclaiming of the dense particles.

If mass does fall into the T_{c-k} , the higher mass M_{tckmax} in the T_{c-k} , causes more infall into the CR. The resulting increase in re-radiated photons causes a decrease in M_{tckmax} . From Eq. (42), the ϵ determines the stable value of M_{tckmax} . This is the observed feedback controlled mechanism required by Merritt and Ferrarese [35] as proposed by the CUM.

C. KR

The start of the KR is caused by a sudden and rapid change in particle type. Almost all the very dense particles (low m_s/m_i) are in the T_{c-k} shell. As r_k increases, the number of dense particles per unit radius decreases, the resident particles's m_s/m_i term is greater, and the M_{tckmax} is much larger than the gravitational effects of particles in the KR upon each other. The observed Keplerian motion of particles in the KR [15, 16] implies the mass M_k within a sphere of radius r_k is approximately M_{tckmax} . Also, since $F_g \propto \tau_f$ of the test particle, the observed Keplerian motion of particles in the KR implies the τ_f of the test particle is nearly constant in the orbit of each test particle.

The m_s/m_i term is causing lighter and denser particles to move out of the KR. The m_s/m_i for hydrogen and the lighter elements is much higher than the massive particles and the m_g is lower. Thus, for hydrogen in the KR the predominant term is the ϵ term which pushes hydrogen and lighter particles out of the KR. The only lighter elements in the KR are gravitationally bound to the denser particles. Black holes are either in or on their way to T_{c-k} . Therefore, Eq. (42) for particles in the KR becomes

$$v_k^2 = r_k \ddot{r}_k + \frac{m_g}{m_i} \frac{GM_{\text{eff}}}{r_k} - K_k, \quad (44)$$

where

$$M_{\text{eff}} \approx M_{\text{tckmax}} - \frac{G_s}{G} \frac{m_s}{m_g} \epsilon, \quad (45)$$

v_k is the v for particles in the KR, and K_k is a relatively small constant accounting for the $\mathbf{a}_o$ term. An upper case K followed by a sub number or letter denotes a constant.

The $\ddot{r}_k$ in Eq. (44) has a value to cause elliptical orbits for the particles which stay in the KR.

As r_k increases, $v_k \rightarrow 0$ in a Keplerian decline. Since differing particles have different m_s/m_l ratios such as stellar material and hydrogen, the minimum v_k for different particle species is reached at different r_k .

The sign of $\mathbf{r}_k \bullet \mathbf{a}_o$ is opposite on opposite sides of the galaxy. Also, the distance between r_k on one side to r_k on the other side is of the order of a few kpc. Compared to the other terms, $\mathbf{r}_k \bullet \mathbf{a}_o$ is small. Therefore, the residual of the $\mathbf{r}_k \bullet \mathbf{a}_o$ from each side of the galaxy is very small. Therefore, the luminosity curve as a function of r_k will be only slightly different on each side of the center. Therefore, averaging the rotation velocity from side to side reduces the effect of relatively small $\mathbf{r}_k \bullet \mathbf{a}_o$ term.

Since the M_{tckmax} is concentrated in a small volume, the KR is spherical. Galaxies are usually inclined relative to our view. Because KR is spherical, the diameter of the KR is measured as the distance along the major axis of the ellipse (from our view).

Since the m_s/m_l particle type differs with radius, this Paper posits that in the high force volume of the KR the m_s/m_l particle types are in strata and the strata's radius depends on ϵ . If each m_s/m_l particle type has a different mass to luminosity ratio, the surface brightness I versus r curve ($I - r$ curve) along the major axis has discontinuities or inflection points. Define r_{ed} (in pc) as the average radius along the $I - r$ curve in each direction from the Source to the first measured discontinuity or inflection point. Therefore, Eq. (25) suggests $r_{\text{sc}} \propto r_{\text{ed}}$ and

$$\epsilon = K_{\text{ed}} r_{\text{ed}}, \quad (46)$$

where K_{ed} is the proportionality constant.

Further, the abrupt change in m_s/m_l particle type occurs in all galaxies. Therefore, r_{ed} may be used to compare parameters among galaxies.

Evaluating Eqs. (42) and (46) at r_{ed} , solving for r_{ed} with the net effect $\mathbf{r}_k \bullet \mathbf{a}_o \approx 0$ when values from each side of the galaxy are averaged yields

$$K_{\text{ed}} = \frac{m_l}{m_s G_s} \left[v^2(r_{\text{ed}}) - \frac{m_g G M(r_{\text{ed}})}{m_l r_{\text{ed}} - r_{\text{ed}} \ddot{r}_{\text{ed}}} \right], \quad (47)$$

where $M(r_{\text{ed}})$ is the mass within a sphere of radius r_{ed} and $v(r_{\text{ed}})$ is the rotation velocity of particles at r_{ed} .

The term in the braces is the difference between the measured $v^2(r_{\text{ed}})$ and the Newtonian expectation.

Substituting Eq. (46) into Eq. (45) yields

$$M_{\text{eff}} = M_{\text{tckmax}} - \frac{G_s}{G} K_{\text{ed}} \frac{m_s}{m_g} r_{\text{ed}} + K_{\text{ieff}}, \quad (48)$$

where K_{ieff} is the intercept of the $M_{\text{eff}} - r_{\text{ed}}$ plot. Theoretically, $K_{\text{ieff}} = 0$ and is a result of measurement error.

1. $M_{\bullet} - r_{\text{ed}}$ data

The kinematic measure of $M_{\bullet}$ uses the relation $r\sigma_v/G$. Therefore, from Eq. (44) $M_{\bullet} \approx M_{\text{eff}}$, where the mass in the KR is relatively much smaller than the mass in the $T_{\text{c-k}}$.

To test Eq. (48), $M_{\bullet}$ data using kinematic methods and distance data for 20 sample galaxies were taken from Merritt and Ferrarese [35]. The data for the 20 sample galaxies are listed in Table I. The data points are the measured values for the galaxies for which the SBH sphere of influence (r_h in Merritt and Ferrarese [35]) has been resolved. The morphology data were obtained from the NED database [78]. The sample includes E, SO, and S type galaxies.

The r_{ed} data were calculated from the distance in Refs. [35] and from the average angular (observed) radius r_a (arcsec). The r_a for each sample galaxy was measured from Hubble Space Telescope (HST) images [79] listed in Table I. A Visual Basic program (see Appendix B) was written to extract the $I - r$ curve and calculate r_a . The selection of each image was based upon being in the V band, upon having a sufficient area of the center of the galaxy in the image so r_a on both sides of the galaxy could be measured, and with sufficient exposure time to have the I value [80] at r_a sufficiently large. The $I - r$ curve obtained from the HST images using the WFPC2 instrument are shown in Fig. 5. NGC 4607 which is listed in Merritt and Ferrarese [35] lacked a satisfactory image. Since the goal of the program is to select a pixel at a discontinuity, image analysis techniques such as neighbor pixel averaging, azimuthal averaging, and combining more than one image into a single data set were avoided. The Visual Basic program has alternate means to deal with peak (center) saturation, bad high pixel values, and bad low pixel values.

The straight line in each $I - r$ curve in Fig. 5 is drawn between the pixels the Visual Basic program selected to calculate r_a . Note, the line is at differing I levels among the galaxies and the slope of the line indicates asymmetry in I value for the calculation of r_a in the same image. Therefore, the selection of r_a is not based on selecting an isophotal value. The r_a is not an isophotal based effective radius.

Figures 6 shows plots of $M_{\bullet}$ versus r_{ed} of the selected galaxies. The straight lines were drawn by (1) identifying the half of the galaxies furthest from the origin, (2) drawing lines from the origin to the identified galaxy points, (3) noting six lines could be matched to the identified galaxies, (4) the galaxies were organized into classifications according to their proximity on the graph to a line, (5) finding the slope and intercept using the least squares method for each classification group of galaxies. The differing slopes of the lines marked by filled diamonds, filled squares, filled triangle, "X", star, and "+" suggest either differing m_s/m_g values, differing K_{ed} values, or differing M_{tckmax} in Eq. (48) among the galaxy classifications. That the data points are on a straight line




FIG. 5: Plots of the $I-r$ profile for the sample galaxies. The straight line is drawn between the chosen pixels. The slope on the line indicates the asymmetry.

TABLE I: Data of the galaxies in Fig. 5.

Galaxy ^a	Morphology	HST		Distance ^b Mpc	r_{ed} ^c pc	$M_{\bullet}$ ^d $10^8 M_{\odot}$	line ^e n
		image	filter				
I1459	E3; LINER	U2BM0102T	F555W	30.3	99.5±14.7	4.6 ± 2.8	2
N0221	cE2	U2E20309T	F555W	0.8	3.7± 0.4	0.039± 0.009	1
N2787	SB(r)0+ LINER	U39S0304R	F555W	7.5	14.6± 3.6	0.41 ^{+0.04} _{-0.05}	2
N3031	SA(s)ab: LINER Sy1	U32L0105T	F547M	3.9	5.5± 1.9	0.68 ^{+0.07} _{-0.13}	3
N3115	S0-	U2J20B04T	F555W	9.8	49.1± 4.8	9.2 ± 3.0	4
N3245	SA(r)ˆ0ˆ	U3MJ1103R	F547M	20.9	35.5±10.1	2.1 ± 0.5	3
N3379	E1: LINER	U2J20F03T	F555W	10.8	51.7± 5.2	1.35 ± 0.73	2
N3608	E2 LINER	U2BM0502T	F555W	23.6	56.5±11.4	1.1 ^{+1.4} _{-0.3}	2
N4258	SAB(s)bc;LINER Sy1.9	U6712101R	F547M	7.2	12.1± 3.5	0.39 ± 0.034	2
N4261	E2-3: LINER	U2I50204T	F547M	33.0	189.4±16.0	5.4 ± 1.2	2
N4342	S0-	U32Q0206T	F555W	16.7	36.9± 8.1	3.3 ^{+1.9} _{-1.1}	3
N4374	E1: LERG LINER	U34K0102T	F547M	18.7	240.2± 9.1	17 ⁺¹² _{-6.7}	3
N4473	E5	U3071503T	F555W	16.1	49.1± 7.8	0.8 ^{+1.0} _{-0.4}	1
N4486	cE0	U2900104T	F547M	16.7	32.1± 8.1	35.7 ±10.2	6
N4564	E6	U3CM7202R	F702W	14.9	26.0± 7.2	0.57 ^{+0.13} _{-0.17}	2
N4649	E2	U2QO0301T	F555W	17.3	52.7± 8.4	20.6 ^{+5.2} _{-10.2}	5
N5128	S0pec	U4100106M	F555W	4.2	33.2± 2.0	2.4 ^{+3.6} _{-1.7}	3
N5845	E	U3070903T	F555W	28.5	167.3±13.8	2.9 ^{+1.7} _{-2.7}	2
N6251	E:LERG Sy2	U2PQ0701T	F555W	104	206.9±50.4	5.9 ± 2.0	2
N7052	E	U2P90108T	F547M	66.1	400.3±32.1	3.7 ^{+2.6} _{-1.5}	1

^a“I” means IC and “N” means NGC..^bThe distance is as stated in the reference for $M_{\bullet}$.^cThe r_{ed} (kpc). The error is ± 0.1 arcsec.^dThe $M_{\bullet}$ was obtained from Merritt and Ferrarese [35].^eThe line is the “n” listed in Table II.




FIG. 6: Plot of the measured mass of the theorized supermassive black hole $M_{\bullet}$ ($10^8 M_{\odot}$) at the center of galaxies versus the distance along the major axis of a HST image to the first discontinuity or inflection point r_{ed} (pc) using the distance stated in Merritt and Ferrarese [35]. The filled diamonds, filled squares, filled triangle, filled circles, open diamond, and open square are data points along a linear $M_{\bullet} - r_{\text{ed}}$ line using data from Table I and correspond to $n = 1$, $n = 2$, $n = 3$, $n = 4$, $n = 5$, and $n = 6$ as listed in Table II, respectively. The error bars (see Table I) were omitted for clarity.

implies the m_s/m_g ratio (particle type), K_{ed} , or M_{tckmax} is discrete for an ϵ value. Since the lines of $n=1$ to $n=3$ (the lines with more than one point) nearly intersect at $(r_{\text{ed}}, M_{\bullet})=(0,0)$ and $M_{\text{tckmax}} > 0$, Eq. (48) implies

$$M_{\text{tckmax}} = K_{\bullet\epsilon}\epsilon = K_{\bullet\epsilon}K_{\text{ed}}r_{\text{ed}}, \quad (49)$$

where $K_{\bullet\epsilon}$ is a proportionality constant. The K_k of Eq. (44) produces a small offset in the intercept of the lines of Eq. (48).

Therefore,

$$M_{\bullet} \approx M_{\text{eff}} = K_{\text{seff}}r_{\text{ed}} + K_{\text{ieff}}, \quad (50)$$

where

$$K_{\text{seff}} = K_{\bullet\epsilon}K_{\text{ed}} - \frac{G_s}{G}K_{\text{ed}}\frac{m_s}{m_g} \quad (51)$$

and K_{seff} is the slope of the $M_{\bullet} - r_{\text{ed}}$ plot.

For a spherically symmetric particle, m_s increases with the square of the radius and m_l increases with the cube of the radius. For a given density, larger particles have lower m_s/m_l . Hydrogen is found in the outer regions of a galaxy and heavier elements are found in the inner regions. Stars larger than our sun are found both inward and outward of us. Therefore, hydrogen has a higher m_s/m_l than iron. Black holes have a much lower m_s/m_l than other stellar particles. Therefore, the largest cross-section area on which F_s acts is on the area of nuclei

TABLE II: Data of the lines in Figures 6.

n^a	line ^b	Corr. ^c	$K_{\text{seff}}^{d,f}$ $10^8 M_{\odot} \text{ kpc}^{-1}$	$K_{\text{ieff}}^{e,f}$ $10^8 M_{\odot}$
1	filled diamonds	0.99+	9.0 ± 0.3	0.11 ± 0.07
2	filled squares	0.92	27 ± 6	0.07 ± 0.8
3	filled triangles	0.99+	70 ± 2	0.1 ± 0.3
4	filled circle	-	187	0
5	open diamond	-	391	0
6	open square	-	1113	0

^aThe integer denoting the place of the line in the order of increasing K_{seff} .

^bThe identification of the line symbol in Figures 6.

^cThe correlation coefficient.

^dThe least squares fit slope of the $M_{\bullet} - r_{\text{ed}}$ lines.

^eThe least squares fit intercept of the $M_{\bullet} - r_{\text{ed}}$ lines.

^fAll lines are calculated as the least squares best fit for points and $(r_{\text{ed}}, M_{\bullet}) = (0,0)$.

rather than on atoms, compounds, or larger assemblies of matter. Gravitationally bound assemblies of matter of differing sizes and the same relative elemental composition will experience the same M_{eff} . Also, assemblies of matter with differing relative elemental composition will have differing RCs. Differing m_s/m_l cause differing v^2 at a given r . For instance, the H_{α} (found near hot stars) and $H\gamma$ RCs are different at small r because the particles are very different. However, in the disk as r increases, the stellar elemental composition is of successively lighter elements. Therefore, the H_{α} and $H\gamma$ RCs become similar at larger r .

The slopes among the lines in Table II obey the relation

$$\log_{10} \left(\frac{S_{\text{lope}}}{U_{\text{nit}}} \right) = K_{\text{sn}}n + K_{\text{in}}, \quad (52)$$

where S_{lope} is the slope of the lines from the parameter relationship; U_{nit} is the units of measure of S_{lope} ; and K_{sn} and K_{in} are the slope and intercept of the linear relation of Eq. (52), respectively. For the $M_{\bullet} - r_{\text{ed}}$ relationship, $S_{\text{lope}} = K_{\text{seff}}$, U_{nit} is $10^8 M_{\odot} \text{ kpc}^{-1}$, $K_{\text{sn}} = 0.41 \pm 0.01$, and $K_{\text{in}} = 0.58 \pm 0.04$ at one standard deviation (1σ). The n is an integer of one to six depending on the line from Table II. The correlation coefficient and F test [81] for Eq. (52) are 0.999 and 0.992, respectively. Note, the lowest slope corresponds to $n=1$ of Eq. (52) and the lines of Figures 6 intersect at a point. Note, S_{lope} is nearly independent of the distance used to calculate the $M_{\bullet}$ and r_{ed} if the same distance is used for both. The values of $\log_{10}(e) \approx 0.434$ is within the 2.3σ of $\log_{10}(K_{\text{sn}})$. Therefore, restating Eq. (52) as a strong Principle of Repetition yields

$$K_{\text{seff}} = K_{\text{se}} e^n, \quad (53)$$

where $K_{\text{se}} = (3.7_{-0.9}^{+0.0}) \times 10^8 M_{\odot} \text{ kpc}^{-1}$ at 1σ .

Combining Eqs. (50) and (53) and setting $K_{\text{iff}} = 0$ yields

$$M_{\bullet} = K_{\text{se}} e^n r_{\text{ed}}. \quad (54)$$

The standard deviation of the χ^2 function are the error bars of $M_\bullet$ listed in Table I. The $\chi^2 = 17$ and the probability value of the χ^2 $P_\chi \approx 60\%$. The data does not invalidate Eq. (54) at the significant 5% level. The χ^2 and P_χ were calculated using only the galaxy data (not the $(r_{\text{ed}}, M_\bullet) = (0, 0)$ point). The $M_\bullet$ is a function of two parameters, r_{ed} and n . The r_{ed} is a parameter of each galaxy, the n is a galaxy classification parameter, and the e implies the Principle of Repetition is applicable to the $M_\bullet$ parameter. Given a set of parameters (n, r_{ed}), a unique value of $M_\bullet$ can be calculated.

A feature of Fig. 6 is the lack of data points in the upper right of the plot (galaxies with relatively high r_{ed} and high $M_\bullet$ values).

D. RR

In the KR as r_k increases to its maximum value (r_{kmax}), the rotation velocity v_{kmin} decreases to its minimum value and the Space and gravitational forces balance. This implies

$$M_{\text{eff}} \rightarrow 0, \quad (55a)$$

$$v_{\text{kmin}}^2 \rightarrow r_{\text{kmax}}(\ddot{r}_{\text{kmax}} - \mathbf{r}_{\text{kmax}} \bullet \mathbf{a}_o). \quad (55b)$$

Therefore, in T_{k-r} the motion of the test particle is primarily governed by the force from external galaxies. Particles such as hydrogen are being ejected outward from the KR with a large radial velocity can cross the T_{k-r} . Particles such as dense stars in nearly circular orbits on a slow inward journey may lack the necessary radial velocity to cross the T_{k-r} . The T_{k-r} is a centrifugal barrier to inward movement of particles. Therefore, external galaxies can have a large influence on the mass distribution and total mass in a galaxy.

If $v_{\text{kmin}} = 0$ in a galaxy, then the CR and KR are decoupled from the RR and SR. The v at $r > r_{\text{kmax}}$ is as if there were no mass inside r_{kmax} and, for non-relativistic calculations, the particles in the RR and SR of an intrinsic galaxy ($\mathbf{a}_o = 0$) are subject to only Newtonian forces.

Two changes in the physics occur to cause the RR. The first change in the first RR (FRR) is to change the particle elemental species which changes m_s/m_l while maintaining a spherical region shape. The second change in the second RR (SRR) is to change the dr increment shape from a spherical shell to a cylindrical shell while continuing the changing m_s/m_l . The resulting equations depend on the v reaching a minimum value ($dv/dr = 0$) of the RC in the KR for the particle species examined.

That shapes of astronomical particle assemblies are either spheroidal or disk is well established. This Paper takes the disk shape as having cylindrical symmetry. The well-known peanut shaped profile of a galaxy bulge is a varying height, cylindrical symmetry with a spheroid core.

1. First RR

The rotation velocity v_r of a particle in the RR is greater than v_{kmin} . If the radius r_{tkr} of a particle's orbit in the T_{k-r} decreases to less than r_{kmax} , the attractive force decreases because M is decreasing. Thus, the effect of ϵ which is spherically symmetric about the center of the galaxy will cause the particle to accelerate to a greater orbit for a given rotation velocity. Therefore, elliptical orbits of particles in the T_{k-r} are inhibited and $\ddot{r}_{\text{tkr}} \approx 0$. Remember one of the simplifying assumptions was the stars are stable in material composition rather than changing m_s/m_l . Changing m_s/m_l would result in a slow inward spiral. Therefore, the radius r_{frr} of the orbits of the particles in the FRR become nearly circular and $\ddot{r}_{\text{frr}} \approx 0$. Thus, the mass in an elemental volume is $D_{\text{frr}} 4\pi r_{\text{frr}}^2 dr$ where D_{frr} is the density of matter at r_{frr} . The D_{frr} is approximately constant because the particles are of the same species. Therefore, the M in Eq. (42) is

$$M_{\text{frr}}(r_{\text{frr}}) = \frac{4}{3} D_{\text{frr}} \pi r_{\text{frr}}^3 + M_{\Delta\text{frr}}, \quad (56)$$

where

$$M_{\Delta\text{frr}} \equiv M_{\text{tkrmax}} - \frac{4}{3} D_{\text{frr}} \pi r_{\text{tkrmax}}^3; \quad (57)$$

$M_{\text{frr}}(r_{\text{frr}})$ is the mass within r_{frr} as a function of r_{frr} ; M_{tkrmax} is the total mass in the CR, T_{c-k} , KR, and T_{k-r} ; and r_{tkrmax} is the radius of the T_{k-r} .

Substituting Eq. (56) into Eq. (42) and taking $\ddot{r} \approx 0$ yields

$$v_{\text{frr}}^2 = \frac{m_g}{m_l} \frac{4}{3} G D_{\text{frr}} \pi r_{\text{frr}}^2 + \frac{m_g G M_{\Delta\text{frr}} - m_s G_s \epsilon}{m_l r_{\text{frr}}} - \mathbf{r}_{\text{frr}} \bullet \mathbf{a}_o(r_{\text{frr}}), \quad (58)$$

where v_{frr} is the rotation velocity of a particle at r_{frr} .

The r_{frr}^{-1} term was small at $r = r_{\text{kmax}}$ and is decreasing as r_{frr} increases. Therefore, the r_{frr}^{-1} term is relatively much smaller than the r_{frr}^2 term. Since the m_s term for hydrogen differs from the m_s term for stellar material, the RCs will differ. By averaging the opposite sides to reduce the $\mathbf{a}_o$ term as before yields

$$v_{\text{frr}}^2 \approx \frac{m_g}{m_l} \frac{4}{3} G D_{\text{frr}} \pi r_{\text{frr}}^2 + K_2, \quad (59a)$$

$$\equiv K_{\text{frrslope}} r_{\text{frrmax}}^2 + K_{\text{frrintercept}}, \quad (59b)$$

where K_{frrslope} and $K_{\text{frrintercept}}$ are the slope and intercept of the v_{frr}^2 - r_{frr}^2 curve.

Figure 7 shows the H I rotation curve of NGC 4321 [9]. It has the rotation velocity plotted well into the bulge. The curved line is in the FRR and is a plot of v_{frr}^2 versus r_{frr} according to Eq. (59).




FIG. 7: Plot of rotation velocity $v^2(10^3 \text{ km}^2 \text{ s}^{-2})$ versus r (kpc) for NGC 4321 [9] using the distance to the galaxy calculated using Cepheid variables to calculate r . A linear curve has been fitted to the SRR. A $v^2 - r^2$ curve has been fitted to the FRR.

2. Second RR

The end of the FRR and the beginning of the SRR is caused by a change in the m_s/m_l factor. The Space force starts to become significant because of m_s/m_l relative to the gravitational force. As the r increases, the particle orbits become rotationally flattened and remain circular [29, pages 723-4]. The spherical symmetry is ended. If either the rotation velocity v_{SRR} in the SRR, the Space force, or the gravitational force changes, the acceleration $\ddot{r}_{\text{SRR}}$ in the SRR will change to restore balance and the circular orbits. Thus, the Principle of Negative Feedback applies and $\ddot{r}_{\text{SRR}} \approx 0$. The orbits of the particles are nearly circular except for changes over a long period in the ϵ ; for changes in the elemental composition and mass of the particles such as by photon emission; and for changes in the $r \bullet a_o$ term. Some of the galaxy's matter, especially the lighter matter with high m_s/m_l , is moving radially. Most of a matter in the SRR is in a stable, radial position. Light material such as hydrogen and heavier elements differ in the mechanics defining their orbits due to the Space force.

The mass in an elemental volume dr of the SRR can be modeled as a cylinder shell of height H_{SRR} [29, page 724] and with density D_{SRR} . The mass in an elemental volume of the SRR is $D_{\text{SRR}}H_{\text{SRR}}2\pi r_{\text{SRR}}dr_{\text{SRR}}$ where r_{SRR} is the radius of a particle's orbit in the SRR. Unlike the inner regions, the m_s/m_l of particles vary more slowly with increasing r_{SRR} . The height profile of the bulge is a measure of the amount of mass versus m_s/m_l particle type. The interplay of the terms of Eq. (42) determine the orbital radius of a particle. Thus, the mass M_{SRR}

within the r_{SRR} is

$$M_{\text{SRR}} = (D_{\text{SRR}}H_{\text{SRR}}2\pi)r_{\text{SRR}}^2 + M_{\Delta r}, \quad (60)$$

where

$$M_{\Delta r} \equiv M_{\text{frrmax}} - (D_{\text{SRR}}H_{\text{SRR}}2\pi)r_{\text{frrmax}}^2; \quad (61)$$

r_{frrmax} is the r to the outer edge of the FRR; and M_{frrmax} is the mass in the CR, T_{c-k} , KR, T_{k-r} , and FRR.

Inserting Eq. (60) into Eq. (42) and with circular orbits yields

$$v_{\text{SRR}}^2 = \left(\frac{m_g}{m_l} G D_{\text{SRR}} H_{\text{SRR}} 2\pi \right) r_{\text{SRR}} + \frac{m_g G M_{\Delta r} - m_s G_s \epsilon}{m_l r_{\text{SRR}}} - r_{\text{SRR}} \bullet a_o. \quad (62)$$

Since the particles in the SRR have larger m_s/m_l and since the r_{SRR}^{-1} term is small at $r = r_{\text{kmax}}$, the r_{SRR}^{-1} term is small compared to the r_{SRR} term. Equation (62) suggest v_{SRR}^2 is nearly independent of ϵ and depends on galaxy parameters $M_{\Delta r}$, D_{SRR} and H_{SRR} . By averaging from side-to-side of the galaxy as before, the $r_{\text{SRR}} \bullet a_o$ term is reduced. Galaxies with other close galaxies in the v plane of rotation, close galaxies with high ϵ values, or several close galaxies asymmetrically arranged, the residual of the side-to-side variation of the $r_{\text{SRR}} \bullet a_o$ term may be relatively significant. Asymmetry in the RC reflects this condition [13]. Therefore,

$$v_{\text{SRR}}^2 = K_{\text{sr}} r_{\text{SRR}} + K_{\text{ir}}, \quad (63)$$

where K_{sr} and K_{ir} are the slope and intercept of the linear $v_{\text{SRR}}^2 - r_{\text{SRR}}$ relationship, respectively. The linear relationship of Eq. (63) for NGC4321 is the straight line plotted in Fig. 7.

To test Eq. (63), galaxies with a distance D_c calculated using Cepheid variables (the “ D_z ” of Freedman et al. [53]) and with a published H I RC which included data points in the SRR were chosen. Figure 8 shows the data points for each RC of the 15 selected galaxies. The straight lines in Fig. 8 are found by fitting a 0.99 or higher correlation coefficient, straight line to the first three data points immediately before the knee of the RC which indicates the transition region T_{r-s} between the RR and SR. NGC 4414 had only two data points before the knee. NGC 3198 and NGC 0224 have several data points between the knee and the straight line of which no three are on a 0.99 correlation straight line.

The data for the 15 selected galaxies are shown in Table III. The morphology type data was obtained from the NED database. The morphology type code t and luminosity class code lc data was obtained from the LEDA database [82]. The number “No.” of data points establishing the SRR, the values of the maximum radius r_{rmax} (kpc) of the last data point on the SRR straight line, the maximum rotation velocity v_{rmax} (km s^{-1}) of




FIG. 8: Plots of the rotation velocity v^2 ($10^3 \text{ km}^2 \text{ s}^{-2}$) versus radius r . The straight lines mark the SSR.

the last data point on the SRR straight line, K_{sr} , and K_{ir} were obtained from the plots of Fig. 8. The “Ref.” lists the references for each RC.

This sample has low surface brightness (LSB), medium surface brightness (MSB), and high surface brightness (HSB) galaxies; galaxies with a range of the 21-cm line width W_{20} at 20 percent of peak of from 120 km s^{-1} to 607 km s^{-1} ; includes LINER, Sy, HII and less active galaxies; galaxies which have excellent and poor agreement between distances D_{tf} calculated using the Tully-

Fisher relation (TF) and D_c ; a D_c range of from 0.75 Mpc to 17.39 Mpc; field and cluster galaxies; and galaxies with rising, flat, and declining RCs in the SR.

The CUM predicts a $v_{\text{frr}}^2 - r_{\text{frr}}^2$ relationship if a galaxy has spherical symmetry in a FRR. At least five of the galaxies in Fig. 8 have data points sufficiently far inward to show the $v_{\text{frr}}^2 - r_{\text{frr}}^2$ relation and don’t. The CUM suggests these galaxies lack a FRR (spherical symmetry).

As r_{srr} increases, m_s/m_l increases and the predominant particles have less mass and decreasing density. De-

TABLE III: Data for the chosen galaxies.

Galaxy	morphology ^a	t ^b	lc ^c	RC ^d	D _c ^e	No.	r _{rmax}	v _{rmax} ^f	K _{sr} ^g	K _{ir} ^h	Ref.
NGC 0224	SA(s)b	3	2	F	0.79	3	10.3±3.7	217±22	5 100± 200	-5 900±11 000	[54]
NGC 0300	SA(s)d	6.9	6	R	2.00	6	3.1±2.2	72±13	1 700± 100	-400± 400	[55]
NGC 0598	SA(s)cd	6	4	R	0.84	5	2.4±1.1	85±12	1 400± 300	4 000± 4 000	[56]
NGC 2403	SAB(s)cd	6	5	F	3.22	3	4.5±0.9	124±10	2 860± 80	2 000± 1 000	[57]
NGC 2841	SA(r)b;LINER Sy	3	1	F	14.07 ⁱ	3	8.3±2.4	324±18	26 400± 700	-8 000± 4 000	[57, 58]
NGC 3031	SA(s)ab;LINER Sy1.8	2.4	2	D	3.63	7	7.4±0.8	243±14	3 900± 200	30 000± 5 000	[59]
NGC 3198	SB(rs)c	5.2	3	F	13.80	4	5.0±1.2	134±15	3 100± 300	2 000± 3 000	[60]
NGC 3319	SB(rs)cd;HII	5.9	3.3	R	13.30	6	8.7±1.7	101± 5	1 300± 100	-1 000± 4 000	[61]
NGC 4258	SAB(s)bc;LINER Sy1.9	4	3.9	D	7.98	3	2.3±0.7	197±26	36 000±10 000	-43 000±30 000	[62]
NGC 4321	SAB(s)bc;LINER HII	4	1	D	15.21	6	5.8±0.8	233±14	5 100± 500	20 000±10 000	[9]
NGC 4414	SA(rs)c? LINER	5.1	3.6	D	17.70	2	6.4±2.9	247±45	6 000	20 000	[63]
NGC 4535	SAB(s)c	5	1.9	D	15.78	3	4.6±1.2	166±12	7 400± 300	-7 000±17 000	[64]
NGC 4548	SBb(rs);LINER Sy	3.1	2	D	16.22	3	10.6±2.9	195±12	3 700± 300	-1 000±17 000	[65]
NGC 5457	SAB(rs)cd	5.9	1	D	6.70	4	8.5±1.3	228±12	4 000± 400	20 000±20 000	[66]
NGC 7331	SA(s)b;LINER	3.9	2	F	14.72	3	2.0±0.6	211±20	11 000± 2 000	23 000±14 000	[67]

^aGalaxy morphological from the NED database.

^bGalaxy morphological type code from the LEDA database.

^cGalaxy luminosity class code from the LEDA database.

^dGalaxy's HI rotation curve type according to slope in the outer SR region. R is rising, F is flat, and D is declining.

^eThe distance D_c (Mpc) to the galaxy from [53] unless otherwise noted.

^fThe maximum rotation velocity v_{rmax} (kms-1), the radius r_{rmax} (kpc) of v_{rmax}, and the No. of points on the straight line from the curves in Fig. 8. The error of r_{rmax} and v_{rmax} is ±5% of the value plus the difference of the chosen data values and the next data point value.

^gLeast squares slope of v_r² - r_r for each galaxy in the SRR of HI rotation curve (km² s⁻² kpc⁻¹).

^hLeast squares intercept of v_r² - r_r for each galaxy in the SRR of HI rotation curve (km² s⁻²).

ⁱThe D_c is from [68].

creasing amount of mass in a given cylindrical shell and the rotation flattening effect causes H_{SRR} to decrease to a disk. This is end of the RR and the beginning of the transition region T_{r-s} between the RR and SR.

3. RR data

Because this change is caused by the change in elemental types which depends on the ϵ of a galaxy, the end of the RR is occurring under similar conditions in all galaxies. Thus, the mass M_{rmax} of the galaxy inside a sphere with a radius of r_{rmax} can be used to compare parameters among galaxies.

Evaluating Eq. (42) at r_{rmax} and rearranging terms yields

$$r_{\text{rmax}} v_{\text{rmax}}^2 = \frac{m_g}{m_l} G M_{\text{rmax}} - \frac{m_s}{m_l} G_s K_{\text{ed}} r_{\text{ed}} + K_{\text{irr}}, \quad (64)$$

where

$$K_{\text{irr}} = r_{\text{rmax}}^2 \ddot{r}_{\text{rmax}} - r_{\text{rmax}} \mathbf{r}_{\text{rmax}} \bullet \mathbf{a}_0. \quad (65)$$

The $r_{\text{rmax}} v_{\text{rmax}}^2$ term is proportional to the effective mass M_{effr} inside a sphere with a radius of r_{rmax} . Since $\ddot{r}_{\text{rmax}} \approx 0$, K_{irr} is a relatively small value. However, in addition to the close-galaxy caveat, a very young galaxy or a galaxy with significant infall of matter such as colliding galaxies can cause a deviation in K_{irr} .

Since r_{rmax} is in the RR and not in T_{r-s} or SR, v_{rmax} can be found in rising, flat and falling RCs. The gas spectral lines such as H_α reflects the motion of a hot star and not hydrogen gas. The elemental content of hot stars changes with r which changes the m_s/m_l term. Therefore, v_{rmax}^2 changes with each star elemental type. The HI RC is monitoring the same particle type in each galaxy from the RR to the SR. Therefore, the HI RC was used to test Eq. (64).

The r_{ed} data listed in Table IV were calculated from D_c in Table III and from the average angular (observed) radius r_a (arcsec). The r_a for each sample galaxy was measured from the Hubble Space Telescope (HST) images listed in Table IV. The Visual Basic program listed in Appendix B was used to extract the $I - r$ curve and calculate r_a . The selection of each image was based upon being in the V band, upon having a sufficient area of the center of the galaxy in the image so r_a on both sides of the galaxy could be measured, and with sufficient exposure time to have the I value at r_a sufficiently large. The $I - r$ curve obtained from the HST images using the WFPC2 instrument are shown in Fig. 9. NGC 2841 had only one side of the galaxy in the image.

The straight line in each $I - r$ curve of Fig. 9 is drawn between the pixels the Visual Basic program selected to calculate r_a . Note, the line is at differing I levels among the galaxies and the slope of the line indicates asymmetry in I value for the calculation of r_a in the same image.

TABLE IV: Data of the galaxies in Fig. 9.

Galaxy ^a	HST image	filter	r_{ed}^b pc	line ^c n m p s
N0224	U2LG0209T	F555W	2.87±0.38	- 6 7 7
N0300	U6713701M	F547M	3.44±0.97	- 2 5 3
N0598	U3MR0106M	F555W	1.50±0.41	- 3 6 5
N2403	U6712504R	F547M	6.61±1.56	- 3 4 4
N2841	U65M7008M	F555W	15.68±6.82	- 5 4 5
N3031	U32L0105T	F547M	5.11±1.76 ^d	3 5 5 6
N3198	U29R1602T	F606W	64.20±6.69	- 1 1 1
N3319	U29R1702T	F606W	19.05±6.45	- 2 4 2
N4258	U6712101R	F547M	13.41±3.87 ^d	2 3 2 4
N4321	U2465107T	F555W	23.66±7.38	- 3 3 4
N4414	U67N8102R	F606W	39.95±8.58	- 3 2 3
N4535	U2782603T	F555W	31.25±7.65	- 2 2 3
N4548	U34L0303T	F555W	32.34±7.87	- 3 3 3
N5457	U6712704R	F547M	5.32±3.25	- 5 5 6
N7331	U41V1802M	F450W	19.23±7.14	- 2 1 4

^a“I” means IC and “N” means NGC.

^bThe r_{ed} was measured as outlined in the text. The error is ± 0.1 arcsec.

^cThe line is the “n” listed in Table II and shown in Fig. 6, the “m” listed in Table V and shown in Fig. 10, the “p” listed in Table VI and shown in Fig. 11, and the “s” listed in Table VII and shown in Fig. 12.

^dThe r_{ed} values differ from Table I because of differing distances used.

TABLE V: Data of the lines in Figures 10.

m^a	line ^b	Corr. ^c	F Test	K_{srr}^d $10^7 \text{ km}^2 \text{ s}^{-2}$	K_{irr}^e $10^4 \text{ kpc km}^2 \text{ s}^{-2}$
1	filled diamonds	ND ^g	ND	0.14	0
2	filled squares	0.99+	0.99+	0.42±0.03	0.3±0.5
3	filled triangles	0.97	0.94	1.1 ±0.1	-0.4±3
4	ND				
5	filled circles	0.97	0.96	5.2 ±0.9	9 ±8
6	open diamond	ND	ND	16.8	0

^aThe integer denoting the place of the line in the order of increasing K_{seff} .

^bThe identification of the line symbol in Figures 10.

^cThe correlation coefficient.

^dThe least squares fit slope of the $r_{\text{max}}v_{\text{max}}^2 - r_{\text{ed}}$ lines.

^eThe least squares fit intercept of the $r_{\text{max}}v_{\text{max}}^2 - r_{\text{ed}}$ lines.

^fThe lines are calculated using $(r_{\text{ed}}, r_{\text{max}}v_{\text{max}}^2) = (0,0)$ as a data point.

^g“ND” means no data because there are only two points (one data point).

Therefore, the selection of r_a is not based on selecting an isophotal value. The r_a is not an isophotal based effective radius.

Figure 10 is a plot of $r_{\text{max}}v_{\text{max}}^2$ versus r_{ed} . Because the sample is small for six galaxy line classifications (four of the six line classifications have less than four galaxies), the $(r_{\text{ed}}, r_{\text{max}}v_{\text{max}}^2) = (0,0)$ point was included in all classifications. Table V lists the calculated values for the lines of Fig. 10.

The lines are straight lines. Therefore, the equation

for each line is

$$r_{\text{max}}v_{\text{max}}^2 = K_{\text{srr}}r_{\text{ed}} + K_{\text{irr}}, \quad (66)$$

where K_{srr} and K_{irr} are the slopes and intercepts of the lines in Fig. 10, respectively.

Since $M_{\text{rmax}} > 0$, the intercepts of the lines in Fig. 10 nearly intersect at $(r_{\text{ed}}, r_{\text{max}}v_{\text{max}}^2) = (0,0)$, and K_{irr} is determined by neighboring galaxies,

$$M_{\text{rmax}} = K_{\text{re}}\epsilon = K_{\text{re}}K_{\text{ed}}r_{\text{ed}}, \quad (67)$$

where K_{re} is the proportionality constant.

The slopes among the lines in Table V obey the relation

$$\log_{10} \left(\frac{K_{\text{srr}}}{\text{km}^2 \text{ s}^{-2}} \right) = K_{\text{srr}}m + K_{\text{irr}}, \quad (68)$$

where $K_{\text{srr}} = 0.40 \pm 0.02$ and $K_{\text{irr}} = 5.79 \pm 0.06$ at 1σ are the least squares fit of the slope and intercept of the linear relation of Eq. (68), respectively. The m is an integer of one to six depending on the line from Table V. The correlation coefficient and F test for Eq. (68) are 0.99 and 0.99, respectively. Note, the lowest slope corresponds to $m=1$ in Eq. (68) and the lines nearly intersect at a point. The K_{srr} is nearly independent of the distance used to calculate the $r_{\text{max}}v_{\text{max}}^2$ and r_{ed} if the same distance is used for both. The values of $\log_{10}(e) \approx 0.434$ is within 1.7σ of K_{srr} .

For the $r_{\text{max}}v_{\text{max}}^2 - r_{\text{ed}}$ relationship,

$$K_{\text{srr}} = \frac{m_g}{m_l} G K_{\text{re}} K_{\text{ed}} - \frac{m_s}{m_l} G_s K_{\text{ed}}. \quad (69)$$

Therefore, restating Eq. (68) as a strong Principle of Repetition yields

$$K_{\text{srr}} = K_{\text{ser}} e^m, \quad (70)$$

where $K_{\text{ser}} = (4.8 \pm 0.9) \times 10^5 \text{ km}^2 \text{ s}^{-2}$ at 1σ .

Combining Eqs. (66) and (70) yields

$$r_{\text{max}}v_{\text{max}}^2 = K_{\text{ser}} e^m r_{\text{ed}}, \quad (71)$$

where $K_{\text{irr}} = 0$.

The standard deviation of the χ^2 function are the error bars of r_{ed} listed in Table IV. The $\chi^2 = 11$ and $P_\chi \approx 70\%$ between Eq. (71) and the measured $r_{\text{max}}v_{\text{max}}^2$ (not including the $(r_{\text{ed}}, r_{\text{max}}v_{\text{max}}^2) = (0,0)$ point). Since P_χ is greater than the 5% significance level, this test does not invalidate Eq. (71). The $r_{\text{max}}v_{\text{max}}^2$ is a function of r_{ed} and m . The r_{ed} is a parameter of each galaxy, the m is a galaxy classification parameter, and the e implies the Principle of Repetition is applicable to the $r_{\text{max}}v_{\text{max}}^2$ parameter, which is proportional to the effective mass at r_{max} .

Since $r_{\text{max}}v_{\text{max}}^2$ is at the outer edge of a m_s/m_l strata, it may be used to compare parameters among galaxies. Posit that r_{max} and v_{max}^2 are related to r_{ed} , also.




FIG. 9: Plots of the $I - r$ profile for the sample galaxies. The straight line is drawn between the chosen pixels. The slope on the line indicates the asymmetry. The horizontal scale for each plot is approximately ten times the r_a .

Figure 11 shows a plot of r_{rmax} versus r_{ed} for the sample galaxies in Tables III and IV. Because the sample is small for seven galaxy integer p classifications (all of the seven p classifications have less than four galaxies), the $(r_{\text{ed}}, r_{\text{rmax}}) = (0, 0)$ point was included in all classifications. The linear relationships are

$$r_{\text{rmax}} = K_{\text{srd}} r_{\text{ed}} + K_{\text{ird}}, \quad (72)$$

where K_{srd} and K_{ird} are the slope and intercept of the

$r_{\text{rmax}} - r_{\text{ed}}$ relationship, respectively, and the respective values are shown in Table VI.

The relationship of the K_{srd} values is

$$\log_{10} K_{\text{srd}} = K_{\text{serd}} p + K_{\text{ierd}}, \quad (73)$$

where $K_{\text{serd}} = 0.28 \pm .02$ and $K_{\text{ierd}} = 1.63 \pm 0.08$ at 1σ .

Note, $\log_{10} 2 \approx 0.3010$. The p is a power of two rather than e . This implies the strong Principle of Change is




FIG. 10: Plot of the measured $r_{\text{max}} v_{\text{max}}^2$ ($10^5 \text{ kpc km}^2 \text{ s}^{-2}$) versus the distance r_{ed} (kpc) along the major axis of a HST image to the first discontinuity or inflection point using the D_c stated in Table IV. The filled diamonds, filled squares, filled triangles, filled circles, and open diamond are data points along a linear $r_{\text{max}} v_{\text{max}}^2 - r_{\text{ed}}$ line using data from Tables III and IV and correspond to $m = 1$, $m = 2$, $m = 3$, $m = 5$, and $m = 6$ as listed in Table V, respectively. The $m=6$ line is drawn between NGC 0224 and $(r_{\text{ed}}, r_{\text{max}} v_{\text{max}}^2) = (0,0)$. The error bars were omitted for clarity.




FIG. 11: Plot of the measured r_{max} (kpc) versus the distance r_{ed} (kpc) along the major axis of a HST image to the first discontinuity or inflection point using the D_c stated in Table IV. The filled diamonds, filled squares, filled triangles, filled circles, open diamonds, open square, and open triangle are data points along a linear $r_{\text{max}} - r_{\text{ed}}$ line using data from Tables III and IV and correspond to $p = 1$, $p = 2$, $p = 3$, $p = 4$, $p = 5$, $p = 6$, and $p = 7$ as listed in Table VI, respectively. The lines are drawn to include the point $(r_{\text{ed}}, r_{\text{max}}) = (0,0)$. The error bars were omitted for clarity.

TABLE VI: Data of the lines in Figures 11.

p^a	line ^b	Corr. ^c	F Test	$K_{\text{srd}}^{d,f}$	$K_{\text{ird}}^{e,f}$ kpc
1	filled diamonds	0.99+	0.99+	76 ± 9	0.2 ± 0.4
2	filled squares	0.99+	0.99+	156 ± 8	0.03 ± 0.2
3	filled triangles	0.98	0.98	310 ± 60	-0.3 ± 1
4	filled circles	0.98	0.98	459 ± 60	0.6 ± 0.8
5	open diamonds	0.96	0.95	1500 ± 300	-0.6 ± 1
6	open square	ND ^g	ND	1620	0
7	open triangle	ND	ND	3590	0

^aThe integer denoting the place of the line in the order of increasing K_{srd} .

^bThe identification of the line symbol in Figures 11.

^cThe correlation coefficient.

^dThe least squares fit slope of the $r_{\text{max}} - r_{\text{ed}}$ lines.

^eThe least squares fit intercept of the $r_{\text{max}} - r_{\text{ed}}$ lines.

^fThe lines are calculated using $(r_{\text{ed}}, r_{\text{max}}) = (0,0)$ as one of the data points.

^g“ND” means no data because there are only two points (one data point).

applicable and

$$K_{\text{srd}} = K_{\text{serdt}} 2^p, \quad (74)$$

where $K_{\text{serdt}} = 36 \pm 8$ at 1 σ .

Combining Eqs. (72) and (74) and setting $K_{\text{ird}} = 0$ yields

$$r_{\text{max}} = K_{\text{serdt}} 2^p r_{\text{ed}}. \quad (75)$$

The standard deviation of the χ^2 function are the error bars of r_{ed} listed in Table IV. The $\chi^2 = 10$ and $P_\chi \approx 76\%$ between Eq. (75) and the measured r_{max} (not including the $(r_{\text{ed}}, r_{\text{max}}) = (0,0)$ point). Since P_χ is greater than the 5% significance level, this test does not invalidate Eq. (75). The r_{max} is a function of r_{ed} and p . The r_{ed} is a parameter of each galaxy, the p is a galaxy classification parameter, and the 2 implies the Principle of Change is applicable to the r_{max} parameter.

Posit

$$v_{\text{max}}^2 = K_{\text{srvd}} r_{\text{ed}} + K_{\text{irvd}}, \quad (76)$$

where K_{srvd} and K_{irvd} are the slope and intercept of the linear relation, respectively.

Using the data from Tables III and IV, the plot of v_{max}^2 versus r_{ed} is shown in Fig. 12 and the data for the lines is listed in Table VII. Because the sample is small for seven galaxy integer s classifications (five of the seven s classifications have less than four galaxies), the $(r_{\text{ed}}, v_{\text{max}}^2) = (0,0)$ point was included in all classifications. The relationship of the slopes of the lines is

$$\log_{10} \left(\frac{K_{\text{srvd}}}{\text{km}^2 \text{ s}^{-2} \text{ kpc}^{-1}} \right) = K_{\text{servd}} s + K_{\text{iervd}}, \quad (77)$$

where $K_{\text{servd}} = 0.31 \pm 0.02$ and $K_{\text{iervd}} = 5.16 \pm 0.07$ at 1 σ .

The strong Principle of Change is applicable and

$$K_{\text{srvd}} = K_{\text{servd}} 2^s, \quad (78)$$




FIG. 12: Plot of the measured v_{rmax}^2 ($\text{km}^2 \text{s}^{-1}$) versus the distance $r_{\epsilon d}$ (kpc) along the major axis of a HST image to the first discontinuity or inflection point using the D_c stated in Table III. The filled diamonds, filled squares, filled triangles, filled circles, open diamond, open triangles, and open square are data points along a linear $v_{rmax}^2 - r_{\epsilon d}$ line using data from Tables III and IV and correspond to $s = 1$, $s = 2$, $s = 3$, $s = 4$, $s = 5$, $s = 6$, and $s = 7$ as listed in Table VII, respectively. The error bars were omitted for clarity.

TABLE VII: Data of the lines in Figures 12.

s^a	s^b	Corr. ^c	F Test	$K_{\text{servd}}^{d f}$ $10^6 \text{ kpc km}^2 \text{s}^{-2}$	$K_{\text{irvd}}^{e f}$ $10^3 \text{ km}^2 \text{s}^{-2}$
1	filled diamonds	ND ^g	ND	0.28	0
2	filled squares	ND	ND	0.54	0
3	filled triangles	0.95	0.92	1.3 ± 0.3	-1 ± 7
4	filled circles	0.99	0.98	2.3 ± 0.2	1 ± 3
5	open diamonds	0.99+	0.99+	6.8 ± 0.1	-1 ± 1
6	open triangles	0.99	0.97	10.6 ± 0.04	0.2 ± 4
7	open square	ND	ND	16	0

^aThe integer denoting the place of the line in the order of increasing K_{servd} .

^bThe identification of the line symbol in Figures 12.

^cThe correlation coefficient.

^dThe least squares fit slope of the $v_{rmax}^2 - r_{\epsilon d}$ lines.

^eThe least squares fit intercept of the $v_{rmax}^2 - r_{\epsilon d}$ lines.

^fThe lines are calculated using $(r_{\epsilon d}, v_{rmax}^2) = (0, 0)$ as one of the data points.

^g“ND” means no data because there are only two points (one data point).

where $K_{\text{servds}} = (1.6 \pm 0.3) \times 10^5 \text{ km}^2 \text{s}^{-2} \text{kpc}^{-1}$ at 1σ .

Combining Eqs. (76) and (78) yields

$$v_{rmax}^2 = K_{\text{servds}} 2^s r_{\epsilon d}, \quad (79)$$

where $K_{\text{irvd}} = 0$.

The standard deviation of the χ^2 function are the error bars of $r_{\epsilon d}$ listed in Table IV. The $\chi^2 = 8.2$ and $P_\chi \approx 88\%$ between Eq. (79) and the measured v_{rmax}^2 (not including the $(r_{\epsilon d}, v_{rmax}^2) = (0, 0)$ point). Since P_χ

is greater than the 5% significance level, this test does not invalidate Eq. (79). The v_{rmax}^2 is a function of $r_{\epsilon d}$ and s . The $r_{\epsilon d}$ is a parameter of each galaxy, the s is a galaxy classification parameter, and the power of two implies the Principle of Change is applicable to the v_{rmax}^2 parameter.

A feature of Figs. 10, 11, and 12 is the lack of data points in the upper right of the plots (galaxies with relatively high $r_{\epsilon d}$ and high r_{rmax} , v_{rmax}^2 , and $r_{rmax} v_{rmax}^2$ values).

E. SR

In the bulge, strong gravitational and Space forces separate the regions into strata by particle m_s/m_l type. In the T_{r-s} , the decreasing height H_{trs} of a galaxy's cylindrical shells results in a lower slope of the plot of the square of the rotation velocity v_{trs}^2 in the T_{r-s} versus radius r_{trs} in the T_{r-s} ($v_{\text{trs}}^2 - r_{\text{trs}}$ curve) than in the RR.

In the SR, the M and ϵ forces are less than in the KR. Therefore, the tendency to separate into separate m_s/m_l regions is less than in the KR. Therefore, the regions are less distinct. The SR could be modeled with these independent regions as rings as is often done to analyze rotation curves. Since the change of m_s/m_l from one ring to the next is small (unlike in the bulge), the form of the fundamental equation of motion for each ring will change little. Also, since the elemental composition of the stars is changing, the border of the rings may be blurred. For purposes herein, to calculate the mass M_s within a sphere with a radius r_s in the SR, the SR is considered as a whole with a smoothly varying rather than stepwise changing m_s/m_l .

The thin disk of the SR is modeled as a cylinder of density D_s and height H_s , then the M_s is

$$M_s = (D_s H_s 2\pi) r_s + M_{\Delta s}, \quad (80)$$

where

$$M_{\Delta s} \equiv M_{\text{trs}} - (D_s H_s 2\pi) r_{rmax} \quad (81)$$

and M_{trs} is the mass inward of the SR.

Because the rotation appears as a rigid body, the projection of the M_s into the RR may be real and not just a mathematical convenience.

Inserting Eq. (80) into Eq. (42) and having circular orbits [29, page 725] yields the rotation velocity v_s in the SR

$$v_s^2 = \left(\frac{m_g G D_s H_s 2\pi}{m_l} \right) r_s + \frac{m_g G M_{\Delta s} - m_s G_s \epsilon}{m_l r_s} - \mathbf{r}_s \bullet \mathbf{a}_0. \quad (82)$$

As in the RR, the r_s^{-1} term is nearly constant. By averaging from side-to-side of the galaxy as before, the $\mathbf{r}_s \bullet \mathbf{a}_0$ term is reduced. However, unlike in the RR,

$D_s H_s < D_{\text{SRR}} H_{\text{SRR}}$. Therefore, $\mathbf{r}_s \bullet \mathbf{a}_o$ is more significant. In cases where there are other close galaxies, close galaxies with moderate ϵ values, several close galaxies asymmetrically arranged, or a close galaxy with a high ϵ , the residual of the side-to-side variation in the $\mathbf{r}_s \bullet \mathbf{a}_o$ term may be relatively significant (see Hodge and Castelaz [13]). Therefore,

$$v_s^2 \approx K_{\text{ss}} r_s + K_{\text{si}} - \mathbf{r}_s \bullet \mathbf{a}_o, \quad (83)$$

where K_{ss} and K_{si} are the slope and intercept of the linear relationship, respectively.

By reasoning similar to the SRR region, the maximum rotation velocity v_{Smax} in the SR is also a linear function of ϵ . Therefore, for the intrinsic galaxy with $\mathbf{r}_s \bullet \mathbf{a}_o \approx 0$

$$v_{\text{Smax}}^2 = K_{\text{seslope}} r_{\text{ed}} + K_{\text{sas}}, \quad (84)$$

where

$$K_{\text{seslope}} = \frac{m_g}{m_l} G K_{\text{es}} - \frac{m_s}{m_l} G_s K_{\text{ed}} \quad (85)$$

and the constant $K_{\text{es}} = M_{\text{Smax}}/r_{\text{ed}}$.

However, unlike the RR, the last rotation velocity data point measured may not be v_{Smax} and will be strongly influenced by $\mathbf{r}_s \bullet \mathbf{a}_o$.

If the impact of the $\mathbf{r}_s \bullet \mathbf{a}_o$ is small, the RC in the SR is linear and rising. Another result of a small $\mathbf{r}_s \bullet \mathbf{a}_o$ is that the difference of rotation velocity at a given radius from one side to the other of a galaxy is small. However, if $\mathbf{r}_s \bullet \mathbf{a}_o$ has a value comparable to the other terms of Eq. (83) and $\nabla^2(\mathbf{r}_s \bullet \mathbf{a}_o) \approx 0$, the $\mathbf{r}_s \bullet \mathbf{a}_o$ term also increases approximately linearly. Therefore, the SR RC will be approximately flat with a slope depending on the relative value of the terms. If $\nabla^2(\mathbf{r}_s \bullet \mathbf{a}_o)$ is significant, the average rotation curve will decline and there will be significant asymmetry in the RC at a given r_s . Since $\mathbf{r}_s \bullet \mathbf{a}_o$ is in the plane of the galaxy, the asymmetry due to neighboring galaxies found by Hodge and Castelaz [13] is in accordance with Eq. (83).

Consider two adjacent ring volumes in the disk with different elemental composition between the rings and with *uniform* elemental composition in each ring. Within each ring, Eq. (44) applies and the RC will be Keplerian. Because of the differing elemental composition, the v_s between the rings obey Eq. (83) and the outer ring will have a greater v_s than the inner ring if $\mathbf{r}_s \bullet \mathbf{a}_o = 0$. However, the differing m_s/m_l is small. Therefore, the detection of the slight decline within each ring may be difficult.

F. DR

The DR is where particles are no longer rotating around the galaxy. The galaxy forces are comparable to the forces from other galaxies. Equation (42) reduces to

$$v_d^2 = r_d \ddot{r}_d - \mathbf{r}_d \bullet \mathbf{a}_{\text{od}}, \quad (86)$$

where

$$\mathbf{a}_{\text{od}} = \sum_{j=1}^{\text{all galaxies}} \frac{m_g G M_j - m_s G_s \epsilon_j}{m_l r_{ij}^3} \mathbf{r}_{ij} + \sum_{k=1}^{\text{Sinks}} \frac{m_g G M_k + m_s G_s \eta_k}{m_l r_{ik}^3} \mathbf{r}_{ik} \quad (87)$$

and the sum is now over all galaxies.

Equation (87) has the form of the Newtonian gravitational force with a anti-gravitational like force, both centered at the center of a galaxy.

IV. OTHER RELATIONSHIPS

A. SAMPLE DATA

The sample of galaxies was drawn from the sample of Freedman et al. [53], who present a sample of 31 spiral galaxies with distances D_c (Mpc) calculated using Cepheid variables (“ D_z ” in Freedman et al. [53]). Since r_{ed} was desired, the sample required each galaxy to have a HST, optical FITS image of the central few kpc of the galaxy. NGC 0925 lacked an adequate HST image and, therefore, was excluded from the sample. NGC 2841 [68] was added to the sample since the D_c determining method was consistent with Freedman et al. [53] and because NGC 2841 is a potential falsifier of the MOND model [5].

The data for the 31 selected galaxies are shown in Tables III, VIII, and IX. The morphology type data were obtained from the NED database. The luminosity class code lc and de Vaucouleurs radius R_{25} data were obtained from the LEDA database. The H I (21 cm) line width W_{20} (km s^{-1}) at 20% of the peak value corrected for inclination data were calculated from “incl” and “w20” parameters from the LEDA database. The m_b = “a magnitude in the visible range” - “extinction” from the NED database. The methods and equations used to calculate the F_p , s , u , w , and x values are described in Section IV B.

The r_{ed} data listed in Table VIII were calculated from D_c listed in Tables III and VIII and from r_a . The r_a for each sample galaxy was measured from the HST images, which were obtained using the WFPC2 instrument, listed in Table VIII. The Visual Basic program listed in Appendix B was used to extract the $I-r$ curve and calculate r_a . The selection of each image was based upon being in the V band, upon having a sufficient area of the center of the galaxy in the image so r_a could be measured, and upon the I value at r_a being sufficiently large. Figure 9 shows the $I-r$ curves for the galaxies with an RC. The remaining $I-r$ curves are shown in Fig. 13. The straight line in each $I-r$ curve of Fig. 13 is drawn between the pixels the Visual Basic program selected to calculate r_a . Note, the line is at differing I levels among the galaxies. The slope of the line indicates asymmetry in I value for

TABLE VIII: Data for the chosen galaxies without rotation curves.

Galaxy	morphology ^a	HST ^b		lc ^c	D_c ^d Mpc	r_{ed} $\times 10^{-3}$ kpc	
		image	filter				
I1613	IB(s)m	U4040201R	F555W	9.8	0.65	1.2±	0.3
I4182	SA(s)m	U6G24404R	F606W	8.1	4.49	5.0±	2.2
N1326A	SB(s)m:	U34L3401O	F555W	7.8	16.14	14.4±	7.8
N1365	(R')SBb(s)b Sy1.8	U2KV010XT	F547M	1	17.95	30.5±	8.7
N1425	SA(rs)b	U29R5P02T	F606W	3.1	21.88	31.9±	10.6
N2090	SA:(rs)b	U29R0W02T	F606W	3.9	11.75	19.8±	5.7
N2541	SA(s)cd LINER	U29R0Z02T	F606W	6	11.22	9.1±	5.4
N3351	SB(r)b;HII Sbrst	U67N3002R	F555W	3	10.00	17.4±	4.9
N3368	SAB(rs)ab;Sy LINER	U6EAN504M	F606W	3	10.52	22.2±	5.1
N3621	SA(s)d	U29R1I02T	F606W	5.8	6.64	10.0±	3.2
N3627	SAB(s)b;LINER Sy2	U29R1K02T	F606W	3	10.05	55.3±	4.9
N4496A	SB(rs)m	U2690102T	F555W	5.9	14.86	14.3±	7.2
N4536	SAB(rs)bc HII	U2DT0601	F555W	1.9	14.93	114.1±	7.2
N4639	SAB(rs)bc Sy1.8	U2NU0202P	F555W	3.8	21.98	135.2±	10.7
N4725	SAB(r)ab pec Sy2	U67N4602R	F606W	2	12.36	28.1±	6.0
N5253	Im pec;HII Sbrst	U2E68101T	F606W	ND	3.15	3.9±	1.5

^aGalaxy morphological from the NED database.^bGalaxy Hubble Space Telescope image from MAST.^cGalaxy data from the LEDA database.^dThe distance D_c (Mpc) to the galaxy from Freedman et al. [53] unless otherwise noted.

TABLE IX: Additional data for the chosen galaxies.

Galaxy	W_{20} ^a km s ⁻¹	R_{25} ^a kpc	F_p ^b $\times 10^6$	m_b mag.	$s u w x$
Galaxies with rotation curves					
N0224	548±10	21.4±1.4	ND ^c	4.26	7 8 - 9
N0300	233±11	5.7±1.6	3.22±0.93	8.95	3 6 5 5
N0598	243± 8	8.1±0.5	ND	6.09	5 8 - 8
N2403	279±18	11.0±1.8	2.98±0.86	8.77	4 6 5 6
N2841	655± 7	15.5±2.0	1.04±0.30	10.09	5 5 3 7
N3031	517± 4	11.8±1.0	0.98±0.28	7.75	6 6 4 8
N3198	343± 7	15.6±2.6	2.89±0.84	10.87	1 3 2 4
N3319	263±13	11.4±1.9	2.01±0.61	11.48	2 4 3 5
N4258	461± 5	19.7±3.3	3.68±1.07	9.10	4 6 4 7
N4321	540±10	16.8±2.5	14.90±4.32	10.00	4 5 5 7
N4414	509± 9	9.1±2.0	1.72±0.50	10.94	3 3 2 5
N4535	423±12	15.9±1.4	6.41±1.86	10.59	3 4 4 6
N4548	449±31	12.4±1.1	0.81±0.23	10.90	3 4 2 5
N5457	502±12	30.1±6.7	3.14±0.91	8.31	6 7 5 9
N7331	538± 7	22.9±3.4	1.00±0.29	10.02	4 5 3 7
Galaxies without rotation curves					
I1613	68± 6	1.5±0.3	ND	9.86	1 5 - 2
I4182	134±11	3.9±0.3	22.80±6.61	13.00	1 5 7 1
N1326A	130±16	4.0±0.9	24.90±7.22	13.77	1 3 6 3
N1365	465±15	28.6±2.5	1.66±0.48	10.32	3 5 3 7
N1425	392±10	19.2±2.5	3.48±1.01	11.87	3 4 3 5
N2090	319±10	9.8±2.9	1.60±0.46	11.99	2 4 3 4
N2541	230±11	9.4±2.4	4.19±1.22	12.09	2 5 5 5
N3351	424± 6	11.0±0.7	1.55±0.45	10.49	4 4 3 5
N3368	439±14	11.9±1.0	32.70±9.48	10.06	4 4 6 6
N3621	306± 6	11.3±2.5	0.05±0.01	9.79	3 5 1 6
N3627	449±10	12.7±1.4	0.70±0.20	9.64	2 3 1 5
N4496A	235± 7	8.2±0.5	33.20±9.63	11.93	2 4 6 5
N4536	394±14	15.7±1.7	19.50±5.66	11.16	1 2 4 3
N4639	420±43	9.7±0.6	26.90±7.80	12.19	1 1 4 2
N4725	510±14	19.7±1.3	19.70±5.71	10.08	4 5 5 6
N5253	101±14	2.2±0.2	0.58±0.173	10.68	1 4 4 4

^aGalaxy data from the LEDA database.^bThe units of F_p are erg cm⁻² s⁻¹ Mpc⁻².^c“ND” means No Data because the Milky Way would be one of the close galaxies.

the calculation of r_a . Therefore, the selection of r_a is not based on selecting an isophotal value. The r_a is not an isophotal based effective radius.

This sample has low surface brightness (LSB), medium surface brightness (MSB), and high surface brightness (HSB) galaxies; galaxies with a range of W_{20} from 68 km s^{-1} to 655 km s^{-1} ; includes LINER, Sy, HII and less active galaxies; a D_c range of from 0.65 Mpc to 21.98 Mpc; field and cluster galaxies; and galaxies with rising, flat, and declining RCs in the disk region.

B. RESULTS

Figure 14 shows a plot of W_{20} versus v_{rmax} . The filled diamonds use the measured values of W_{20} and v_{rmax} of the galaxies with an RC. The line is a plot of,

$$\frac{W_{20}}{\text{km s}^{-1}} = K_1 \left(\frac{v_{\text{rmax}}}{\text{km s}^{-1}} \right)^{K_2} (\pm 12\%), \quad (88)$$

where $K_1 = 7.3$ and $K_2 = 0.78$.

Equation (79) relates r_{ed} and v_{rmax} . The s for each galaxy in Table IX without a RC was chosen to provide the data point closest to the line of Eq. (88).

The uncertainties of r_{ed} listed in Table I were used to calculate the standard deviation of the χ^2 function. The $\chi^2 = 8$ and $P_\chi \approx 86\%$ between Eq. (88) and the measured W_{20} . Since P_χ is greater than the 5% significance level, this test does not invalidate Eq. (88). The W_{20} is a function of v_{rmax} that is a function of r_{ed} and s . The r_{ed} is a parameter of each galaxy, the s is a galaxy classification parameter, and the “2” implies the Principle of Change is applicable to the W_{20} parameter.

The W_{20} is used by the Tully-Fisher relationship [69, 70] to calculate total absolute magnitude M_β (mag.) in band β

$$M_\beta = K_{4\beta} + K_{5\beta} \log(W_{20}), \quad (89)$$

where $K_{4\beta}$ and $K_{5\beta}$ are constants that depend on the band under consideration. Therefore, M_β is a function of s and r_{ed} .

The correlation coefficients of W_{20} versus r_{rmax} and $r_{\text{rmax}}v_{\text{rmax}}^2$, which is proportional to the effective mass within a sphere of radius r_{rmax} , are 0.38 and 0.80, respectively.

Figure 15 is a plot of R_{25} versus r_{ed} . Each line in Fig. 15 is a different “classification”. Because four of the eight classifications had less than four galaxies, the $(r_{\text{ed}}, R_{25}) = (0,0)$ point was included in all classifications. Table X lists the calculated values for the lines of Fig. 15.

The lines of Fig. 15 are straight lines. Therefore, the equation for each line is

$$R_{25} = K_{\text{sr}25} r_{\text{ed}} + K_{\text{ir}25}, \quad (90)$$

where $K_{\text{sr}25}$ and $K_{\text{ir}25}$ are the slopes and intercepts of the lines in Fig. 15, respectively.

TABLE X: Data of the lines in Figures 15.

u^a	line ^b	Corr. ^c	F Test	$K_{\text{sr}25}^{d,f}$	$K_{\text{ir}25}^{e,f}$
1	filled diamond	ND ^g	ND	71	0
2	filled square	ND	ND	138	0
3	filled triangles	0.99+	0.99+	233 $\pm$ 9	1 $\pm$ 1
4	filled circles	0.95	0.87	490 $\pm$ 60	1 $\pm$ 1
5	open diamonds	0.95	0.87	800 $\pm$ 100	1 $\pm$ 2
6	open squares	0.97	0.95	1400 $\pm$ 200	1 $\pm$ 2
7	open triangles	0.99+	0.99+	5680 $\pm$ 80	0.2 $\pm$ 0.3
8	open circle	ND	ND	7450	0

^aThe integer denoting the place of the line in the order of increasing $K_{\text{sr}25}$.

^bThe identification of the line symbol in Figures 15.

^cThe correlation coefficient.

^dThe least squares fit slope of the $R_{25} - r_{\text{ed}}$ lines.

^eThe least squares fit intercept of the $R_{25} - r_{\text{ed}}$ lines.

^fThe lines are calculated using $(r_{\text{ed}}, R_{25}) = (0,0)$ as a data point.

^g“ND” means no data because there are only two points (one data point).

Since $K_{\text{ir}25} \approx 0$, the slopes among the lines in Table X obey the relation

$$\log_{10} \left(\frac{R_{25}}{r_{\text{ed}}} \right) = K_{\text{su}} u + K_{\text{iu}}, \quad (91)$$

where $K_{\text{su}} = 0.30 \pm 0.02$ and $K_{\text{iu}} = 1.52 \pm 0.08$ at 1σ are the least squares fit of the slope and intercept of the linear relation of Eq. (91), respectively. The u is an integer of one to eight depending on the line from Table X. The correlation coefficient and F test for Eq. (91) are 0.99 and 0.98, respectively. Note, the lowest slope corresponds to $u = 1$ in Eq. (91). The R_{25}/r_{ed} is nearly independent of the distance used to calculate the R_{25} and r_{ed} if the same distance is used for both. The value of $\log_{10}(2) \approx 0.301$ is well within 1σ of K_{su} .

Therefore, restating Eq. (91) as a strong Principle of Change and combining Eqs. (90) and (91) yields

$$R_{25} = K_{\text{Rr}} 2^u r_{\text{ed}}, \quad (92)$$

where $K_{\text{Rr}} = (32 \pm 7)$ at 1σ and $K_{\text{ir}25} = 0$. Note the lack of data points in the upper right of Fig. 15.

The error bars of r_{ed} listed in Tables IV and VIII were used to calculate the standard deviation of the χ^2 function. The $\chi^2 = 17$ and $P_\chi \approx 96\%$ between Eq. (92) and the measured R_{25} [not including the $(r_{\text{ed}}, R_{25}) = (0,0)$ point]. Since P_χ is greater than the 5% significance level, this test does not invalidate Eq. (92). The R_{25} is a function of r_{ed} and u . The r_{ed} is a parameter of each galaxy, the u is a galaxy classification parameter, and the “2” implies the Principle of Change is applicable to the R_{25} parameter.

If ϵ is the total energy entering a galaxy and luminosity $(10^{-0.4 M_j})$ is the total energy emitted from a galaxy, then $\epsilon \propto 10^{-0.4 M_j}$ in a stable galaxy, where M_j is the total B band absolute magnitude of the j^{th} galaxy and B band luminosity is a fixed proportion of the total luminosity. Tully and Fisher [69] originally suggested luminosity is proportional to its effective mass of the galaxy. Then the




FIG. 13: Plots of the $I - r$ profile for the sample galaxies without a rotation curve. The straight line is drawn between the chosen pixels. The slope on the line indicates the asymmetry.

force F_p ($\text{erg cm}^{-2} \text{s}^{-1} \text{Mpc}^{-2}$) exerted by the mass and ρ of the neighboring galaxies on the target galaxy [71] along the polar axis is

$$F_p = K_{fz} \sum_{j=1}^n 10^{-0.4 M_j} \frac{1}{R_j^3} Z_j, \quad (93)$$

where K_{fz} is the proportionality constant, the sum is over the closest galaxies, n is the number of galaxies included

in the calculation, $\mathbf{R}_j$ is the vector distance from the target galaxy to the j^{th} close galaxy, and Z_j is component of $\mathbf{R}_j$ along the target galaxy's polar axis. In the calculation herein, $K_{fz} = 1$ and $n = 10$.

The calculation followed Hodge and Castelaz [71]. IC 1613, NGC 0224, and NGC 0300 were omitted from the calculation since the Milky Way is one of the closest galaxies and the M of the Milky Way is uncertain. The result of the calculation[83] for 28 of the sample galaxies.




FIG. 14: Plot of W_{20} versus v_{max} . The filled diamonds use the measured values of W_{20} and v_{max} . The line is a plot of Eq. (88).




FIG. 15: Plot of the measured R_{25} (kpc) versus the distance r_{ed} (kpc) along the major axis of a HST image to the first discontinuity or inflection point using the D_c stated in Table III and VIII. The filled diamond, filled square, filled triangles, filled circles, open diamonds, open squares, open triangles, and open circle are data points along a linear $R_{25} - r_{\text{ed}}$ line using data from Table IX and correspond to $u = 1$, $u = 2$, $u = 3$, $u = 4$, $u = 5$, $u = 6$, $u = 7$, and $u = 8$ as listed in Table X, respectively.

The F_p error was calculated using 0.2 mag error for M_j and a 5% error for distance.

Figure 16 is a plot of F_p versus r_{ed} . Each line in Fig. 16 is a different “classification”. The $(r_{\text{ed}}, F_p) = (0,0)$ point was included in all classifications. Table XI lists the calculated values for the lines of Fig. 16.




FIG. 16: Plot of the measured F_p ($\text{erg cm}^{-2} \text{s}^{-1} \text{Mpc}^{-2}$) versus the distance r_{ed} (kpc) along the major axis of a HST image to the first discontinuity or inflection point using the D_c stated in Tables III and VIII. The filled diamonds, filled squares, filled triangles, filled circles, open diamonds, open squares, and open triangle are data points along a linear $F_p - r_{\text{ed}}$ line using data from Table IX and correspond to $w = 1$, $w = 2$, $w = 3$, $w = 4$, $w = 5$, $w = 6$, and $w = 7$ as listed in Table XI, respectively.

TABLE XI: Data of the lines in Figures 16.

w^a	line ^b	Corr. ^c	F Test	$K_{\text{sf}}^{d f}$	$K_{\text{iw}}^{e f}$
1	filled diamonds	0.99	0.99	13.2 ± 0.5	-0.04 ± 0.03
2	filled squares	0.94	0.89	42 ± 3	0.7 ± 0.7
3	filled triangles	0.93	0.86	97 ± 1	-0.07 ± 0.04
4	filled circles	0.99	0.99	185 ± 6	0.4 ± 0.3
5	open diamonds	0.99	0.98	680 ± 30	-0.5 ± 0.6
6	open squares	0.93	0.91	1570 ± 90	3 ± 2
7	open triangle	ND ^g	ND	4570	0

^aThe integer denoting the value of w in Eq. 95.

^bThe identification of the line symbol in Fig. 16.

^cThe correlation coefficient.

^dThe least squares fit slope of the lines in Fig. 16. The units of K_{sf} are $\times 10^6 \text{ erg cm}^{-2} \text{s}^{-1} \text{Mpc}^{-2} \text{kpc}^{-1}$.

^eThe least squares fit intercept of the lines in Fig. 16. The units of K_{iw} are $\times 10^6 \text{ erg cm}^{-2} \text{s}^{-1} \text{Mpc}^{-2}$.

^fThe lines are calculated using $(r_{\text{ed}}, F_p) = (0,0)$ as a data point.

^g“ND” means no data because there are only two points (one data point).

The lines of Fig. 16 are straight lines. Therefore, the equation for each line is

$$F_p = K_{\text{sf}} r_{\text{ed}} + K_{\text{iw}}, \quad (94)$$

where K_{sf} and K_{iw} are the slopes and intercepts of the lines in Fig. 16, respectively.

Unlike the previous r_{ed} relationships, K_{iw} deviates slightly from zero. However, the slopes among the lines

in Table XI obey the relation

$$\log_{10} \left(\frac{K_{\text{sf}}}{\times 10^6 \text{ erg cm}^{-2} \text{ s}^{-1} \text{ Mpc}^{-2} \text{ kpc}^{-1}} \right) = K_{\text{sw}} w + K_{\text{iw}}, \quad (95)$$

where $K_{\text{sw}} = 0.41 \pm 0.01$ and $K_{\text{iw}} = 0.73 \pm 0.05$ at 1σ are the least squares fit of the slope and intercept of the linear relation of Eq. (95), respectively. The w is an integer of one to seven depending on the line from Table XI. The correlation coefficient and F test for Eq. (95) are 0.998 and 0.996, respectively. Note, the lowest slope corresponds to $w = 1$ in Eq. (95). The value of $\log_{10}(e) \approx 0.43$ is within 2σ of K_{sw} .

Therefore, restating Eq. (95) as a strong Principle of Repetition and combining Eqs. (94) and (95) yields

$$F_{\text{p}} = K_{\text{Fr}} e^w r_{\text{ed}}, \quad (96)$$

where $K_{\text{Fr}} = (4.5 \pm 0.8) \times 10^6 \text{ erg cm}^{-2} \text{ s}^{-1} \text{ Mpc}^{-2} \text{ kpc}^{-1}$ at 1σ and $K_{\text{iw}} = 0$.

The error bars of F_{p} listed in Tables IX were used to calculate the standard deviation of the χ^2 function. The $\chi^2 = 27$ and $P_{\chi} \approx 34\%$ between Eq. (96) and the measured F_{p} [not including the $(r_{\text{ed}}, F_{\text{p}}) = (0, 0)$ point]. Since P_{χ} is greater than the 5% significance level, this test does not invalidate Eq. (96). The F_{p} is a function of r_{ed} and w . The r_{ed} is a parameter of each galaxy, the w is a galaxy classification parameter, and the “e” implies the Principle of Repetition is applicable to the F_{p} parameter.

Since $\epsilon = K_{\text{ed}} r_{\text{ed}}$,

$$r_{\text{ed}} = \frac{\epsilon}{K_{\text{ed}}} = \frac{F_{\text{p}}}{K_{\text{Fr}} e^w}, \quad (97)$$

where K_{ed} is the proportionality constant.

Since $\mathbf{Z}_j$ may have any value, the continuous factors of the r_{ed} relationships may be due, in part, to $\mathbf{F}_{\text{p}}$.

Figure 17 is a plot of galaxy luminosity L versus r_{ed} . The L value is calculated as

$$L = 10^{-0.4 M_{\text{V}}}, \quad (98)$$

where

$$M_{\text{V}} = m_{\text{b}} - 25 + 5 \log_{10} D_{\text{c}}, \quad (99)$$

where the “-25” and “5” in the equation are because the unit of D_{c} is Mpc.

Each line in Fig. 17 is a different “classification”. Because six of the nine classifications had less than four galaxies, the $(r_{\text{ed}}, L) = (0, 0)$ point was included in all classifications. Table XII lists the calculated values for the lines of Fig. 17.

The lines of Fig. 17 are straight lines. Therefore, the equation for each line is

$$L = K_{\text{sLr}} r_{\text{ed}} + K_{\text{iLr}}, \quad (100)$$

where K_{sLr} and K_{iLr} are the slopes and intercepts of the lines in Fig. 17, respectively.




FIG. 17: Plot of the measured L ($\text{erg cm}^{-2} \text{ s}^{-1}$) versus the distance r_{ed} (kpc) along the major axis of a HST image to the first discontinuity or inflection point using the D_{c} stated in Tables III and VIII. The filled diamond, filled squares, filled triangles, filled circles, open diamonds, open squares, open triangles, open circles, and “X” are data points along a linear $L - r_{\text{ed}}$ line using data from Table IX and correspond to $x = 1$, $x = 2$, $x = 3$, $x = 4$, $x = 5$, $x = 6$, $x = 7$, $x = 8$, and $x = 9$ as listed in Table XII, respectively.

TABLE XII: Data of the lines in Figures 17.

x^a	line ^b	Corr. ^c	F Test	$K_{\text{sLr}}^{d f}$	$K_{\text{iLr}}^{e f}$
1	filled diamond	ND ^g	ND	26	0.0
2	filled squares	0.99+	0.99+	48 ± 0	0.0 ± 0.0
3	filled triangles	0.99+	0.99+	68 ± 1	-0.1 ± 0.2
4	filled circles	0.99+	0.99+	133 ± 5	-0.1 ± 0.2
5	open diamonds	0.97	0.92	285 ± 20	0.1 ± 0.7
6	open squares	0.99+	0.99+	474 ± 18	0.2 ± 0.4
7	open triangles	0.97	0.94	851 ± 100	2 ± 2
8	open circles	0.99+	0.99+	2100 ± 100	-0.2 ± 0.3
9	X	0.99+	0.99+	4000 ± 200	0.3 ± 0.6

^aThe integer denoting the place of the line in the order of increasing K_{sLr} .

^bThe identification of the line symbol in Figures 17.

^cThe correlation coefficient.

^dThe least squares fit slope ($\text{erg cm}^{-2} \text{ s}^{-1} \text{ kpc}$) of the $L - r_{\text{ed}}$ lines.

^eThe least squares fit intercept ($\text{erg cm}^{-2} \text{ s}^{-1}$) of the $L - r_{\text{ed}}$ lines.

^fThe lines are calculated using $(r_{\text{ed}}, L) = (0, 0)$ as a data point.

^g“ND” means no data because there are only two points (one data point).

Since $K_{\text{iLr}} \approx 0$, the slopes among the lines in Table XII obey the relation

$$\log_{10} \left(\frac{K_{\text{sLr}}}{\text{erg cm}^{-2} \text{ s}^{-1} \text{ kpc}} \right) = K_{\text{sx}} x + K_{\text{ix}}, \quad (101)$$

where $K_{\text{sx}} = 0.274 \pm 0.007$ and $K_{\text{ix}} = 1.08 \pm 0.04$ at 1σ are the least squares fit of the slope and intercept of the linear relation of Eq. (101), respectively. The x is an integer of one to nine depending on the line from Table XII. The

correlation coefficient and F test for Eq. (101) are 0.99 and 0.99, respectively. Note, the lowest slope corresponds to $x = 1$ in Eq. (101). The L/r_{ed} is not independent of the distance used to calculate the L and r_{ed} . The value of $\log_{10}(2) \approx 0.301$ is within 4σ of K_{sx} . There are only three galaxies on the $x = 1$ and $x = 2$ lines. If the $x = 1$ and $x = 2$ lines are ignored, the $K_{\text{sx}} = 0.29 \pm 0.03$ that is within 1σ of $\log_{10}(2)$.

Therefore, restating Eq. (101) as a strong Principle of Change and combining Eqs. (100) and (101) yields

$$L = K_{\text{Lr}} 2^x r_{\text{ed}}, \quad (102)$$

where $K_{\text{Lr}} = (8.9 \pm 2.0) \text{ erg cm}^{-2} \text{ s}^{-1} \text{ kpc}$ at 1σ and $K_{\text{ir25}} = 0$. Note the lack of data points in the upper right of Fig. 17.

The error bars of r_{ed} listed in Tables IV and VIII were used to calculate the standard deviation of the χ^2 function. The $\chi^2 = 30$ and $P_\chi \approx 44\%$ between Eq. (102) and the measured L [not including the $(r_{\text{ed}}, L) = (0, 0)$ point]. NGC 4639 had a $\chi^2 = 18$. Since P_χ is greater than the 5% significance level, this test does not invalidate Eq. (102). The L is a function of r_{ed} and x . The r_{ed} is a parameter of each galaxy, the x is a galaxy classification parameter, and the “2” implies the Principle of Change is applicable to the L parameter.

The F_p is calculated from the magnitude, which is derived from ϵ , and the vector position of other Sources. Since F_p and M are independent variables, r_{ed} of a galaxy depends on the M_j , hence ϵ , of all galaxies. Also, other galaxy parameters are related to r_{ed} . This suggests the ϵ and the position in Euclidian space of all the Sources (center of galaxies) completely determines the parameters of a target galaxy.

V. THE MANY LINES IN THE PARAMETER RELATIONS WITH r_{ed} ARE NOT RANDOM.

The many lines in each plot in Figs. 6, 10, 11, 12, 15, 16, and 17 may suggest the relationships portrayed may be just random data points. The null hypothesis to be tested is “the data points are indeed random points”. This null hypothesis may be tested by following the procedure used in discovering the Eqs. (54), (71), (75), (79), (92), (96), and (102) as follows: (1) Generate 15 sets of two random numbers between zero and one. Call one of the i^{th} set the independent variable (X_i) and the other the dependant variable (Y_i), where i varies from one to 15. (2) The equation to be tested is

$$Y_{\text{calci}} = K_1 2^{t_i} X_i, \quad (103)$$

where K_1 is the proportionality constant and t_i is the integer classification for the i^{th} set. (3) A trial consists of generating the random numbers for the 15 sets and subjecting the sets to the slope, correlation coefficient, and χ^2 calculations and tests the galaxy samples passed. (4) Calculate the K_1 value for each trial as the minimum

value of $K_1 = Y_i/(2^{t_i} X_i)$ with an $X_i > 0.4$. (5) Calculate the t_i value for each set of (X_i, Y_i) values as

$$t_i = \text{ROUND} \left[\log_2 \left(\frac{Y_i}{K_1 X_i} \right) \right], \quad (104)$$

where “ROUND” means to round the value in the braces to the nearest integer. (6) If any one of the $t_i > 7$, the trial failed. (7) For calculating the slopes and correlation coefficients, if any subset lacks a sufficient number of values, that subset will be ignored. (8) If the correlation coefficient of any t_i subset of (X_i, Y_i) values, including the $(X_0, Y_0) = (0, 0)$ point, is less than 0.95, the trial failed. (9) If the linear slope of the relation between the t_i and slope, including the $(X_0, Y_0) = (0, 0)$ point, of the t_i subsets $\neq (0.30 \pm 0.03)$, the trial failed. (10) If the correlation coefficient of the line in (9) above < 0.95 , the trial failed. (11) Calculate the Y_{calci} according to Eq. (103). If the $P_\chi < 0.05$ between the Y_i and Y_{calci} for the 15 sets, the trial failed. (12) maintain a count of the number N_π of sets of (X_i, Y_i) with $X_i^2 + Y_i^2 \leq 1$ and the number N_{XltY} of sets of (X_i, Y_i) with $X_i < Y_i$. (13) Redo the trial with another 15 sets of random numbers for 30,000 trials.

The results are: (1) The $N_\pi/(15 \times 30,000) = 0.78528 \approx \pi/4$ and $N_{\text{XltY}}/(15 \times 30,000) = 0.49999$. Therefore, the random number generator performed satisfactorily. (2) 22 of the 30,000 trials passed (0.07%), the remainder of the trials failed. Therefore, the null hypothesis is rejected with a 99+% confidence.

VI. DISCUSSION

Space has a physical presence, a density, and an ability to exert a force on the α of hods (matter) without being matter. The Space in this Paper looks like the ether with added properties. The Michelson-Morley experiments [72] applied to the velocity of photons. To achieve the results of Michelson-Morley’s experiments, photons must have zero surface area presented to their direction of travel. Hence, photons must be hods arranged in a column. However, recent re-examination of the Michelson-Morley data [73] and more accurate measurements [74] suggests a preferred reference frame may exist. The CUM suggests these findings indicate ripples on the hods of photons provide a dragging force on the photon. Therefore, in addition to testing the ether model, Lorentzian relativity, and Einstein’s special relativity, the Michelson-Morley type experiment may be testing the ρ , $\nabla\rho$, and $d(\nabla\rho)/dt$ in our solar system. The ratio of the diameter of the earth’s orbit to the distance from the center of the Galaxy suggests the Michelson-Morley experiment may require greater precision and appropriate timing to test the ρ , $\nabla\rho$ and $d(\nabla\rho)/dt$.

Ferrarese [39] discovered a linear relationship between circular velocity v_c beyond R_{25} and bulge velocity dispersion σ_c (see her Fig. 1). Baes et al. [75] expanded on the data. NGC 0598 was a clear outlier. Galaxies with σ_c

less than 70 km s^{-1} (v_c less than 150 km s^{-1}) also deviate from the linear relation. Also, galaxies with significant non-circular motion in the HI gas such as NGC 3031 were omitted in Ferrarese [39]. The v_c for NGC 0598 used in Ferrarese [39] was 135 km s^{-1} , which is the highest data point of a rising RC [56]. Most of the other curves Ferrarese [39] used were flat. For flat RCs $v_c \approx v_{rmax}$. If the $v_c = v_{rmax} = 85 \text{ km s}^{-1}$ is used by Ferrarese [39] for NGC 0598, NGC 0598 will fit on her plotted line. Thus, the range of $v_c - \sigma_c$ relation may extend to $27 \text{ km s}^{-1} < \sigma_c < 350 \text{ km s}^{-1}$. NGC 3031 has a declining RC, $R_{25} \approx 11 \text{ arcmin}$ (from the LEDA database), $v_c(R_{25}) \approx 190 \text{ km s}^{-1}$ [59], and $\sigma_c = 174 \pm 17 \text{ km s}^{-1}$ [35]. NGC 3031 is not on the plotted line of Ferrarese [39]. However, if $v_c = v_{rmax} = 243 \pm 14 \text{ km s}^{-1}$ is used, NGC 3031 does fit on the plotted line.

Multiple slopes of the C_a versus E_a ($\text{erg cm}^{-2} \text{ s}^{-1}$) relation from Hodge and Castelaz [71] (see Hodge and Castelaz [71] for the definition and calculation of C_a and E_a) also obey Eq. (52). Separate the CET1 zone into two zones. The first zone (CET1a) has NGC 3621 and NGC 5253 which have low E_a and a slope of $-2.28 \times 10^{-2} \text{ erg}^{-2} \text{ cm}^2 \text{ s}^1$. The second zone (CET1b) has NGC 3031 and NGC 2090 which have higher E_a and a slope of $8.87 \times 10^{-4} \text{ erg}^{-1} \text{ cm}^2 \text{ s}^1$. The slope values were arranged in ascending order with CET1b, CET2, CET3, and CET4 corresponding to $n=1, n=2, n=3$, and $n=4$, respectively. For the C_a versus E_a relationship, U_{nit} is $\text{erg}^{-1} \text{ cm}^2 \text{ s}^1$, $K_{sn} = 0.585 \pm 0.005$ and $K_{in} = -0.54 \pm 0.02$ at 1σ . The correlation coefficient and F test for Eq. (52) are 0.99 and 0.99, respectively.

The linear potential model (LPM) explored by Mannheim and Kmetko [67, and references therein] is derived from relativistic considerations. The LPM proposes a gravitational potential $V(r)$ of the form $V(r) = -\beta/r + \gamma r/2$ where β and γ are constants. The potentials in the CUM are proportional to r^{-1} . RCs with $v_{srr}^2 \propto r_{srr}$ or $v_{frr}^2 \propto r_{frr}^2$ are derived by geometric and particle type variations after the force of the M_{kmax} is balanced by the Space force. However, both models conclude the intrinsic v_{srr} and v_s are a linear function of r . Hence, both predict rising rotation curves. The LPM derives flat curves by mass distribution but ultimately has rising curves. If LPM is to derive declining curves, it is by the influence of other galaxies which have a linear r potential. The LPM does not predict the relations with r_{ed} or $v_{frr}^2 \propto r_{frr}^2$.

The DM model (DMM) places large amounts of unknown, non-luminous matter in a halo around a galaxy to explain rising rotation curves. Whereas, the DMM requires the DM term to become larger for rising rotation curves, CUM requires the $\mathbf{r} \bullet \mathbf{a}_o$ term to become smaller. The disagreement between current CDM theory and observation is most evident in galaxies with low surface brightness (LSB) such as NGC 0300 and rising rotation curves such as NGC 0300, NGC 0598, and NGC 3319. These galaxies fit in the CUM. In addition, the DMM would consider v_{rmax} as merely another point on the rotation curve. The DMM suggests mass is infalling

from outside the galaxy. The CUM suggests matter is outflowing as gas and inflows as denser particles. The DMM is not really a predictive model but rather a parameterization of the difference between observation and Newtonian expectation. Because of the number of parameters in the DMM, it is very difficult to falsify. However, the large number of parameters makes the DMM cumbersome and, therefore, replaceable.

Padmanabhan and Choudhury [76, and references therein] explored the possibility of a scalar field acting as dark matter. The CUM suggests the massless (mass density = 0) scalar ρ field at small scales causes an intrinsic rising RC. The DMM would require a large amount of DM. The ρ field acts like large amounts of DM at galaxy diameter scales. Therefore, the CUM suggests the amount of DM in the universe is zero. Also, this Paper suggest the ρ field causes Newtonian kinematic measurements to considerably underestimate mass in a galaxy. For example, Eq. (45) with $M_{eff} = M_\bullet$ suggests the larger M_{tckmax} is the more accurate measure of the amount of mass in the central region of galaxies.

The CUM's $(m_g M_g - m_s \epsilon)/r$ term has the form of a subtractive change in acceleration. The CUM suggests this causes the species of matter to change with radius which causes the appearance of a gravitational acceleration change with radius. In the MOND model [5], the Newtonian gravitational acceleration is measured in our solar system, which is in the low r end of the SR. The MOND data comparison is usually in the far RR and SR regions. Hence, the existence of a differing ϵ/r or an apparent modified gravitational acceleration is expected by the CUM model. NGC2841 is a candidate to falsify MOND. A distance greater than 19 Mpc, compared to the measured D_c of 14.1 Mpc [68], is needed for MOND [5]. However, NGC2841 data fits the CUM. Further, given a relatively short radius range of the MOND application, the ϵ/r change may be approximated as a constant. Also, since we are near the v_{rmax} in the Galaxy and since the CUM holds the relationship of $v_{rmax}^2 - \epsilon$ as a universal relationship, the MOND proposition that the modified gravitational acceleration is a universal constant follows. If in the SR there is another acceleration from $\mathbf{r} \bullet \mathbf{a}_o$ to balance the ϵ decline, then the MOND's flat rotation curve can result. MOND also predicts that if the rotation curve is flat, it is axis symmetric. Flat and declining rotation curves are asymmetric [13, and references therein]. Rising rotation curves are more symmetric. Therefore, the CUM reduces to MOND where $\Delta \mathbf{r} \bullet \mathbf{a}_o \neq 0$ and $\nabla^2(\mathbf{r} \bullet \mathbf{F}_o) = 0$ in the SR.

The model that places a SBH at the center of galaxies says nothing about the RC outside the KR. Also, this model of the center of galaxies seems to be invalidated by observations of gas near the center of the Galaxy [29, page 595][30], by the apparent inactivity of the SBH [31, 32, 33, 34], by the observed relations with other galaxy parameters [16, 23, 37, 39, 44, 45, 46, 47, 48, 49], and by the apparent need for a feedback control mechanism [35].

The DMM adds several parameters per galaxy. Both

the LPM and MOND add one parameter to Newtonian mechanics and another mildly varying parameter to achieve their fits and have been applied to only close galaxies. The CUM adds one parameter (ϵ) per galaxy and one term per m_s species to Newtonian dynamics, but needs other galaxies and their distances to further explain a galaxy's dynamics. The ϵ of other galaxies modify the equations to allow flat and declining RCs and to allow asymmetries in RCs. Neither the LPM nor MOND address the relationships outside the RR and SR. The DMM, LPM, MOND, and SBH galaxy models either are invalidated by or fail to explain the data examined herein.

I consider it very attractive that the CUM, at this early stage of model development, has brought together heretofore apparently unrelated observations and has discovered new galaxy parameter relations.

VII. CONCLUSION

A changing universe model (CUM) was developed in accordance with a new set of Fundamental Principles. The CUM posits the stuff of our universe is continually erupting into our universe from sources at the center of galaxies. The stuff of our universe is matter and a matterless scalar whose gradient exerts a force on the cross section of matter. An energy continuity equation (Section II A) was derived. Equations describing the forces of gravity (Section II B 3) and Space (Section II B 1) were derived. A description of the structure of particles (Section II C) was presented to provide an explanation of the phenomena observed at the Galaxy center. An explanation of Galaxy structure was presented. The CUM qualitatively suggests: (1) That our universe is flat. (2) The characteristic of the right angle is orthogonality. (3) That total kinetic energy equals total potential energy in our universe. (4) That $\nabla_m \rho$, α , τ , and ι are constants of the universe. (5) That Space is causally connected and coherent. (6) Why there are three, and only three, Space dimensions. (7) Why there are three, and only three, families of particles.

To test the proposition, a single equation was derived using the CUM that calculates the rotation velocity v of particles along the major axis of spiral galaxies. The equation adds one parameter for each mature galaxy and one parameter for each particle (atom nucleus and smaller) species to Newtonian dynamics. The equation is used to correlate the measured mass $M_\bullet$ of the theorized, central supermassive black hole (SBH) with other galaxy parameters and to describe the rotation velocity curve (RC) of spiral galaxies. The application of the CUM suggests galaxies are divided into regions. The CUM suggests the intrinsic rotation curve in the disk region is rising and suggests flat and declining disk rotation curves are caused by neighboring galaxies. Each region of a galaxy has its own physics. The single equation, through the relative value of terms, reduces to a description of the physics of the RC of each region. Four new parameters

were defined: (1) the radius r_{ed} to a discontinuity in the surface brightness versus radius curve, (2) the rotation velocity v_{rmax} at the maximum extent of the rapidly rising region of the RC, (3) the radius r_{rmax} to the v_{rmax} , and (4) the Source strength ϵ of each galaxy.

In a sample of 20 galaxies with the radius of influence of the central black hole resolved, this Paper predicted a linear $M_\bullet - r_{ed}$ relation and found $M_\bullet = K_{se} r_{ed} e^n$, where $K_{se} = (3.7^{+0.0}_{-0.9}) \times 10^8 M_\odot \text{ kpc}^{-1}$, $e \approx 2.718$ (the natural log base), and n is an integer from one to six depending on the galaxy. Since the sample included E, SO, and S galaxies, the derived equation blends smoothly across the galaxy morphologies of the sample.

In a sample of 15 spiral galaxies with H I RCs and Cepheid based distances, this Paper predicted a linear $r_{rmax} v_{rmax}^2 - r_{ed}$ relation and found $r_{rmax} v_{rmax}^2 = K_{ser} r_{ed} e^m$, where $K_{ser} = (4.8 \pm 0.9) \times 10^5 \text{ km}^2 \text{ s}^{-2}$ and m is an integer from one to six depending on the galaxy. Also, this Paper predicted a linear $r_{rmax} - r_{ed}$ relation and found $r_{rmax} = K_{serdt} r_{ed} 2^p$, where $K_{serdt} = 36 \pm 8$ and p is an integer from one to seven depending on the galaxy. Also, this Paper predicted a linear $v_{rmax}^2 - r_{ed}$ relation and found $v_{rmax}^2 = K_{servds} r_{ed} 2^s$, where $K_{servds} = (1.6 \pm 0.3) \times 10^5 \text{ km}^2 \text{ s}^{-2} \text{ kpc}^{-1}$ and s is an integer from one to seven depending on the galaxy. Further, for each of the four linear relationships, linear relationships between the respective parameter and r_{ed} for each of the integer classifications were also found. The CUM predicts the square of the rotation velocity is linearly related to r in a portion of the RR and is linearly related to r^2 in another portion of the RR.

For a sample of 31 spiral galaxies with a wide range of characteristics, the H I (21 cm) line width W_{20} (km s^{-1}) at 20% of the peak value corrected for inclination has been found to be tightly related to v_{rmax} by $W_{20}/\text{kms}^{-1} = K_1 (v_{rmax}/\text{kms}^{-1})^{K_2}$, where $K_1 = 7.3$, $K_2 = 0.78$, and the uncertainty in W_{20} is $\pm 12\%$ with $P_\chi \approx 86\%$. Since W_{20} is related to the total absolute magnitude M_β (mag.) in band β by the Tully-Fisher relationship [69, 70], M_β is also related to r_{ed} . The de Vaucouleurs radius R_{25} (kpc) is proportional to r_{ed} by $R_{25} = K_{Rr} 2^u r_{ed}$ with $P_\chi \approx 96\%$, where $K_{Rr} = (32 \pm 7)$ at 1σ and u is an integer from one to eight depending on the galaxy. For 28 of the 31 sample galaxies, the force F_p of the ϵ , which is proportional to the luminosity, of neighboring galaxies directed along the polar axis of the target galaxy is related to r_{ed} by $F_p = K_{Fr} e^w r_{ed}$ with $P_\chi \approx 34\%$, where $K_{Fr} = (4.5 \pm 0.8) \times 10^6 \text{ erg cm}^{-2} \text{ s}^{-1} \text{ Mpc}^{-2} \text{ kpc}^{-1}$ at 1σ and w is an integer from one to seven depending on the galaxy. This suggests the continuous factors of the r_{ed} relationships are due, in part, to F_p . For the 31 sample galaxies, the luminosity L in the visible band (from the NED database) is proportional to r_{ed} by $L = K_{Lr} 2^x r_{ed}$ with $P_\chi \approx 44\%$, where $K_{Lr} = (8.9 \pm 2.0) \text{ erg cm}^{-2} \text{ s}^{-1} \text{ kpc}$ at 1σ and x is an integer from one to nine depending on the galaxy.

The r_{ed} is related to galaxy dynamics parameters, luminosity parameters, and an isophotal radius parameter

by the Principle of Change. Also, r_{cd} is related to F_p and effective mass parameters by the Principle of Repetition.

The CUM is not invalidated by the data presented herein.

Acknowledgments

This research has made use of the NASA/IPAC Extragalactic Database (NED) which is operated by the Jet Propulsion Laboratory, California Institute of Technology, under contract with the National Aeronautics and Space Administration.

This research has made use of the LEDA database (<http://leda.univ-lyon1.fr>).

This research has made use of data obtained from the Multimission Archive at the Space Telescope Science Institute (MAST). STScI is operated by the Association of Universities for Research in Astronomy, Inc., under NASA contract NAS5-26555. Support for MAST for non-HST data is provided by the NASA Office of Space Science via grant NAG5-7584 and by other grants and contracts.

This research has made use of the NASA's Skyview facility located at NASA Goddard Space Flight Center.

I acknowledge and appreciate the financial support of Cameron Hodge, Stanley, New York, and Maynard Clark, Apollo Beach, Florida, while I was working on this project.

APPENDIX A: PRINCIPLES

The Principal of Fundamental Principles is that a Fundamental Principle, and the models developed from it, is a meaningful and useful principle that applies to all levels of physical systems. To be meaningful is to be able to be used to describe and predict the outcome of experiments. To be useful is to be able to be used to cause desired outcomes. An outcome of an experiment includes placing bounds on what (parameter or event), where (position coordinates), and when (time coordinate). The level of a physical system refers to the size of the domain of applicability over which a set of physical theories applies such as galaxies versus atoms and Newtonian versus special relativity.

Corollary I is if a candidate to be a Fundamental Principle is found to not apply in some system then it is not a Fundamental Principle. Corollary II is if a principle is found in all of physical systems, then it is likely to apply to larger and smaller levels and to new concepts.

The Principle of Superposition, the Correspondence Principle, and Principle of Minimum Potential Energy are such Fundamental Principles. The Correspondence Principle is an interpolation of the Fundamental Principles. Corollary II is an extrapolation of the Fundamental Principle.

A “scientific model” (theory) is derived from the transcendent idea of the Fundamental Principle and is applicable to a defined domain. Since life and social systems are physical systems, by Corollary II and the Cosmological Principle, the transcendent idea of the Fundamental Principles must also apply to life and to social systems. The more fundamental scientific models have larger domains. A proposal becomes a scientific model when a deduction from it can be found to be factual (is observed). This concept does not require a candidate model to include “falsifiable predictions” and does not invalidate a scientific model because of the existence of falsifying observations. For instance, Newtonian dynamics is a valid scientific model. Observations in the domain including relative velocities approaching the speed of light falsify Newtonian dynamics. However, this only limits the Newtonian dynamics domain. Religious ideology, models based on belief, and philosophy models may be scientific models provided they are useful and meaningful with restricted what, where, and when bounds. To survive, a scientific model must compete for attention. The concept of a scientific model survives because the human mind is limited in the ability to maintain a catalog of the nearly infinite number of possible observation. Scientific models with empty or more limited domains have little usefulness and meaningfulness.

The Copernican Principle, which states we are not privileged observers of the universe, is accepted. In quantum mechanics our presence may change the outcome of experiments. However, this is true for any observer. If, in some set of observations, we appear privileged, the privilege must be incorporated in the model. For example, we are in a galaxy disk and are close to a sun. We are in a highly unique and privileged area. Just because we are carbon based does not imply all intelligent life is carbon based.

The Anthropic Principle is accepted to the extent that what is observed must have been created and have evolved to the present. What is observed must be able to be observed. Note this statement of the Anthropic Principle omits the requirement that it depend on an assumption of “life” and “intelligence” since life and intelligence are inadequately defined. The existence of life, social systems, and intelligence are observations of our universe and, therefore, must be able to exist. An unobserved parameter may or may not be able to be observed. Therefore, the negative model candidates are not useful.

Some argue [77, and references therein] the Anthropic Principle cannot be part of science since it cannot yield falsifiable predictions. In this statement of Fundamental Principles, what is observed can be used to define the domain of scientific models.

The Cosmological Principle is accepted to the extent that the physics must be the same at all positions in the universe. Therefore, the theories must explain the conditions at all positions in the universe. For example, the physics that states all objects fall to earth was found to be limited when Galileo noted moons rotating around

Jupiter. Newton's restatement of physics was more cosmological. However, the Cosmological Principle is not extended to imply the universe is isotropic and homogeneous. Near a star is a unique position. The physics must explain other positions. The physical theories must explain any isotropies and anisotropies.

The Principle of Change states that each system is built of previous systems by a minimum step change. When the size of a structure of an object becomes limited, a new structure comprising a combination of existing elements can occur. Change is a building on existing structures. Destruction of objects into its parts does not foster the Principle of Change. Since the minimum step Change is a split from one to two, a corollary (strong Principle of Change) is that whatever exists will split into two parts. The two parts may be a continuation of what was plus a new system or may be what was incorporated into the two new systems. Particles became a hydrogen atom followed by evolution of other atoms. Atoms became molecules. A model that requires a large step where there are possible intervening steps is not observed and is forbidden.

A corollary of the Principle of Change is that all things have a beginning.

The Principle of Repetition states that there are two ways to repeat a Change: (1) If conditions allow an observable change, then the change will occur again under similar conditions. (2) The repeated Changes have a common cause (reproduction). A corollary is that if two systems have the same observable results, then similar conditions exist or the systems were reproduced. A strong statement of the Principle of Repetition is that the amount of increase of a parameter by the Principle of Repetition depends on the size of the parameter. Destruction of objects to have "room" for "the new" is a Repetition because the only objects that can be built from the pieces are a Repetition of objects.

The Principle of Negative Feedback states that any system with a relatively narrow parameter relationships must evolve from a broader system and must have a negative feedback loop to maintain the narrow parameters and achieve balance between the Change and Repetition processes. Otherwise, the system is unstable and transitory. The effects of the unstable system will cease to exist without consequential fallout or permanent change. We observe objects that have limited size. So, there is a limiting force or negative feedback condition controlling the size of each object. So too must black holes have a size limitation and a negative feedback condition.

The Principle of Minimum Potential Energy can be stated as a Principle of Minimum Action, which states the path of least energy expenditure will be followed during the change from one state to another. Combining this with the Principle of Negative Feedback yields the conclusion that the feedback energy equations are linear, of the first time derivative, and of the first time integral. In an equation none, one, or two of these elements may be zero. Since a higher power feedback equation would

result in the same controlled output and consume more energy, the minimum equations are sufficient. Therefore, the assumption of "linear" in formulating descriptive, negative feedback energy equations is sufficient.

The Principle of Geometric Rules states that the observed geometric relationships apply in all levels of systems. Hence, the conservation of energy/mass must be related to geometric rules we observe in our universe. Hence, π = circumference / radius in two dimensions must be the same number in three dimensions. π is not only a universal constant in our universe, it is a constant in all the universes. The division by two is another universal concept. The division by two for each dimension into equal angles yields the right angle.

APPENDIX B: VISUAL BASIC 5.0 PROGRAM FOR FINDING r_a .

This program starts with the appropriate information extracted from the HST FITS file and the operator has input the required additional data.

```
Option Explicit
Const Pi As Single = 3.141592654
'% integer
'& long integer
'! single precision
'# double precision
'$ string
Dim i%, j%, x%, y%, k&, l&, m&, p%
Dim d!(802, 802), dmax!, Xmax!, Ymax!
Dim XL!, YL!, RAp#, DECp#, Dist#, PA!
Dim Ap#, b0#, cp#, Xc#, DECO#
Dim XLv&, YLv&, XLvLast%, YLvLast%, RA0#
Dim crVal1#, crVal2#, crPix1#, crPix2#
Dim NAXIS1%, NAXIS2%, ORIENTAT!, CD1_1#
Dim dmaxI!, XmaxI!, YmaxI!, CD1_2#, CD2_1#
Dim VegaZP#, ExpTime#, Area#, CD2_2#
Dim Instrument$, Filter$, IData!(2270)
Dim AData#(2270), Sb!, TSlope!, R1p!
Dim Sb0!, CountL%, CountH%, R1n!, Rea1!
Dim Rea2!, Sxy#, Sx#, Sx2#, Sy#
'crVal1, crVal2, crPix1, crPix2, ExpTime
'Instrument, Filter,
'NAXIS1, NAXIS2, ORIENTAT, CD1_1, CD1_2
'CD2_1, CD2_2
'are as defined in the HST FITSFILE header.
'VegaZP is the zero point of the surface
'brightness calculation, input by operator
'd(X,Y) is the data number (DN)
'of the pixel
'at the X and Y image coordinate.
'Since the galaxy may be asymmetric,
'd(X,Y) is not averaged or
're-calculated statistically.

Private Sub cmdDoOne_Click()
```

```

    PA = CInt(Text1.Text) 'position
' angle of galaxy, operator input
    Area = Abs(CD1_1 * CD1_2) * 3600 _
* 3600
'find center of gal. by highest pixel
' value close to input coordinates
    Xmax = Val(Text2.Text)
    Ymax = Val(Text3.Text)
    dmax = d(Xmax, Ymax)
DoAgain2:
    For x = Xmax - 20 To Xmax + 20
    For y = Ymax - 20 To Ymax + 20
        If x > 0 And y > 0 And x < _
NAXIS1 And y < NAXIS2 Then
            If d(x, y) > dmax Then
                dmax = d(x, y)
                Xmax = x
                Ymax = y
                GoTo DoAgain2
            End If
        End If
    Next
    Next
    Text2.Text = CStr(Xmax)
    Text3.Text = CStr(Ymax)
    PA = PA - ORIENTAT
' PA is now angle from X axis
'counterclockwise on image and Y axis
'is up from bottom
DoAgain:
    If PA >= 180 Then      'not matter
' for tan function
        PA = PA - 180
        GoTo DoAgain
    End If
    If PA < 0 Then
        PA = PA + 180
        GoTo DoAgain
    End If
    If PA > 89.3 And PA < 90 Then
' TAN(90) is problem
        PA = 89.3
    End If
    If PA < 90.7 And PA > 90 Then
        PA = 90.7
    End If
'record general information in Data1
' which is an Excel file
    Data1.Refresh
    Data1.Recordset.MoveFirst

    Data1.Recordset.Edit
    Data1.Recordset.Fields("NAME").Value_
= Left(File1.filename, _
Len(File1.filename) - 4)
    Data1.Recordset.Update
    Data1.Recordset.MoveNext
    Data1.Recordset.Edit

```

```

    Data1.Recordset.Fields("NAME").Value _
= Label2.Caption + " deg."
    Data1.Recordset.Update
    Data1.Recordset.MoveNext
    Data1.Recordset.Edit
    Data1.Recordset.Fields("NAME").Value_
= Label3.Caption + " deg"
    Data1.Recordset.Update
    Data1.Recordset.MoveNext
    Data1.Recordset.Edit
    Data1.Recordset.Fields("NAME").Value_
= "IMAGE PA: " + Format(PA, "##0.#")
    Data1.Recordset.Update
    Data1.Recordset.MoveNext
    Data1.Recordset.Edit
    Data1.Recordset.Fields("NAME").Value _
= "CNT: " + Format(d(CInt(Text2.Text), _
CInt(Text3.Text))), "#####")
    Data1.Recordset.Update
    Data1.Recordset.MoveNext
    Data1.Recordset.Edit
    Data1.Recordset.Fields("NAME")_
.Value _
= "CNT X: " + Format(Val(Text2.Text)_
, "###")
    Data1.Recordset.Update
    Data1.Recordset.MoveNext
    Data1.Recordset.Edit
    Data1.Recordset.Fields("NAME")_
.Value _
= "CNT Y: " + Format(Val(Text3.Text)_
, "###")
    Data1.Recordset.Update
    Data1.Recordset.MoveNext
    Data1.Recordset.Edit
    Data1.Recordset.Fields("NAME").Value _
= "INSTRUMENT: " + Instrument
    Data1.Recordset.Update
    Data1.Recordset.MoveNext
    Data1.Recordset.Edit
    Data1.Recordset.Fields("NAME").Value_
= "FILTER: " + Filter
    Data1.Recordset.Update
    Data1.Recordset.MoveNext
    Data1.Recordset.Edit
    Data1.Recordset.Fields("NAME").Value_
= "VegaZP:" + CStr(VegaZP)
    Data1.Recordset.Update
    Data1.Recordset.MoveNext
    Data1.Recordset.Edit
    Data1.Recordset.Fields("NAME").Value _
= "EXPOSURE TIME:" + CStr(ExpTime)_
    Data1.Recordset.Update
'initialize
    Data1.Recordset.MoveFirst
    For i = 1 To 2270
        IData(i) = 0
        AData(i) = 0
    Next i

```

```

        Data1.Recordset.Edit
        Data1.Recordset.Fields("INDEX")_
.Value = 0
        Data1.Recordset.Fields("X"_
).Value = 0
        Data1.Recordset.Fields("Y")_
.Value = 0
        Data1.Recordset.Fields("DEGREES")_
.Value = 0
        Data1.Recordset.Fields("VALUE")_
.Value = 0
        Data1.Recordset.Update
        Data1.Recordset.MoveNext
    Next
    Data1.Recordset.MoveFirst
'coordinates of gal. center
    DECO = crVal1 + ((Val(Text2.Text) _
- crPix1) * CD1_1) + ((Val(Text3.Text)_
- crPix2) * CD1_2)
    RA0 = crVal2 + ((Val(Text2.Text)_
- crPix1) * CD2_1) + ((Val(Text3.Text)_
- crPix2) * CD2_2)
'a change of location of center on
'the image
    If Val(Text2.Text) > 0 Then
        crPix1 = Val(Text2.Text)
    End If
    If Val(Text3.Text) > 0 Then
        crPix2 = Val(Text3.Text)
    End If
    p = 0
    If PA < 45 Or PA > 135 Then
        For XL = 1 To NAXIS1 - 1 _
Step 0.25
            XLv = Int(XL)
            YLv = crPix2 + ((XL - crPix1)_
* (Tan(PA * Pi / 180)))
            If YLv > 1 And YLv < NAXIS2 _
- 1 Then
                Record
            End If
        Next
    Else
        For YL = 1 To NAXIS2 - 1 _
Step 0.25
            YLv = Int(YL)
            XLv = crPix1 + ((YL - crPix2)_
* (Tan((90 - PA) * Pi / 180)))
            If XLv > 1 And XLv < NAXIS1_
- 1 Then
                Record
            End If
        Next
    End If
'record last record
    Data1.Recordset.Edit
    Data1.Recordset.Fields("X").Value_

```

```

= CSng(NAXIS1)
        Data1.Recordset.Fields("Y").Value_
= CSng(NAXIS2)
        Data1.Recordset.Fields("VALUE")_
.Value = 0
        Data1.Recordset.Fields("DEGREES")_
.Value = 0
        Data1.Recordset.Update
        Data1.Recordset.Close

    Rea
        Label4.Caption = "DONE: " _
+ File1.filename
End Sub

Sub Record()
    If XLv = XLvLast And YLv _
= YLvLast Then 'prevent
'duplication of pixels
        GoTo NotDo
    End If
    XLvLast = XLv
    YLvLast = YLv
    p = p + 1
    IData(p) = d(XLv, YLv)
'calculate the Right Assension and
' DECLination of the pixel
    RAp = RA0 + ((crPix1 - XLv) * CD1_1_
) + ((crPix2 - YLv) * CD1_2)
    DECp = DECO + ((crPix1 - XLv) _
* CD2_1) + ((crPix2 - YLv) * CD2_2)
    Ap = RAp - RA0 'left in degrees
' for troubleshoot ease
    b0 = 90 - DECO
    cp = 90 - DECp
'calculate the cos(angle from the
'center to the pixel)
    Dist = (Sin(b0 * Pi / 180) * _
Sin(cp * Pi / 180) * Cos(Ap * Pi / 180))_
+ (Cos(b0 * Pi / 180) _
* Cos(cp * Pi / 180))

'calculate arccos of Dist
    If Abs(Dist) > 0.9999999999999995_
Then
        Dist = 0
    Else
        AData(p) = (Atn(-Dist / _
Sqr(-Dist * Dist + 1)) + 2 * Atn(1))_
* 180 / Pi
    End If

'dist >0 unless changed
    If XLv < crPix1 And YLv <=_
crPix2 Then_
        AData(p) = -AData(p)
    End If
    If XLv >= crPix1 And YLv <_

```

```

crPix2 Then_
    AData(p) = -AData(p)
End If
If PA > 90 Then
    AData(p) = -AData(p)
End If
'record DN pixel values
Data1.Recordset.Edit
Data1.Recordset.Fields("INDEX")_
.Value = CSng(p)
Data1.Recordset.Fields("X")_
.Value = CSng(XLv)
Data1.Recordset.Fields("Y")_
.Value = CSng(YLv)
Data1.Recordset.Fields("VALUE")_
.Value = IData(p)
Data1.Recordset.Fields("DEGREES")_
.Value = AData(p)
Data1.Recordset.Update
Data1.Recordset.MoveNext
NotDo:
End Sub

Function Slope2(A As Long) As Double
    If A = 0 Then
        Sxy = 0
        Sx = 0
        Sy = 0
        Sx2 = 0
        l = 0
    Else
        Sxy = Sxy + (IData(A)
* AData(A))
        Sx = Sx + AData(A)
        Sy = Sy + IData(A)
        Sx2 = Sx2 + (AData(m + k)
* AData(m + k))
        l = l + 1
        If Abs((l * Sx2) - (Sx * Sx))_
< 0.000000001 Then
            Slope2 = 0
        Else
            Slope2 = ((l * Sxy) _
- (Sx * Sy)) / ((l * Sx2) - (Sx * Sx))
        End If
    End If
End Function

Sub Rea()
'recalculates and records Rea data

'find index number (m) of the gal.
'center, the center already chosen
redom:
    IData(0) = 0
    For k = 1 To p
        If IData(0) < IData(k) And _
IData(k) <= dmax Then
            IData(0) = IData(k)
            AData(0) = AData(k)
            m = k
        End If
    Next
    For k = 1 To p
        If k <= m And AData(k) > 0 Then
            AData(k) = -AData(k)
        End If
        If k > m And AData(k) < 0 Then
            AData(k) = -AData(k)
        End If
    Next
're-record data as surface brightness
'and angular data in arc seconds
Data1.Refresh
Data1.Recordset.MoveFirst
    For k = 1 To p
        Data1.Recordset.Edit
        Data1.Recordset.Fields("INDEX")_
.Value = CSng(k)
        If IData(k) < 1 Then
            Data1
.Recordset.Fields("VALUE")_
.Value = CSng(25)
        Else
            Data1
.Recordset.Fields("VALUE")_
.Value = CSng(VegaZP - 2.5 * _
(Log(CDbl(IData(k)) / ExpTime / Area)_
/ Log(10#)))
        End If
        Data1
.Recordset.Fields("DEGREES")_
.Value = CSng(3600 * AData(k))
        Data1.Recordset.Update
        Data1.Recordset.MoveNext
    Next
Data1.Recordset.Close

'pos going
'find starting k
    k = 0
    Do While IData(m + k) > 0.95 _
* IData(m)
        k = k + 1
    Loop
    If IData(m + k) < 0.8 * IData(m_
) Then k = -1
    CountH = 0
    TSlope = -10000000
    Sb = -20000000
redoSlopeP3:
    Sb0 = Slope2(0) 'reset slope
' numbers
    For j = 1 To 2 'initialize
' slope
        k = k + 1

```

```

        If IData(m + k) < _
IData(m + k - 1) Or k = 0 Then
            Sb0 = Slope2(m + k)
        Else
            j = j - 1
        End If
    Next
    CountL = 0
    Do While CountL < 10 And _
IData(m + k) > 50 And (m + k + 2) > 1
        k = k + 1
        If IData(m + k) > 5 Then
'for false data pixel
            Sb = Slope2(m + k)
            If Sb0 < Sb Then
                CountL = CountL + 1
                CountH = 0
                Sb0 = Sb
            Else
                CountH = CountH + 1
                CountL = 0
                Sb0 = Sb
            End If
        Else
            IData(m + k) = 60
        End If
    Loop
    Rip = m + k - CountL

'neg going
'find starting k
    k = 0
    Do While IData(m + k) > 0.95 _
* IData(m)
        k = k - 1
    Loop
    If IData(m + k) < 0.8 * IData(m) _
Then k = 1
    CountH = 0
    TSlope = 10000000
    Sb = 20000000
redoSlopen3:
    Sb0 = Slope2(0) 'reset slope
' numbers
    For j = 1 To 2 'initialize
' slope
        k = k - 1
        If IData(m + k) < _
IData(m + k + 1) Or k = 0 Then
            Sb0 = Slope2(m + k)
        Else
            j = j - 1
        End If
    Next
    CountH = 0
    Do While CountH < 10 And _
IData(m + k) > 50 And (m + k - 2) > 1
        k = k - 1

```

```

        If IData(m + k) > 5 Then
'for false data pixel
            Sb = Slope2(m + k)
            If Sb0 < Sb Then
                CountL = CountL + 1
                CountH = 0
                Sb0 = Sb
            Else
                CountH = CountH + 1
                CountL = 0
                Sb0 = Sb
            End If
        Else
            IData(m + k) = 60
        End If
    Loop
    Rin = m + k + CountH

    Rea1 = Abs(AData(R1p) - AData(R1n))
    If Abs(IData(R1p + 2)) < 10 Then
        Rea1 = Abs(AData(m) - _
AData(R1n)) * 2
        IData(R1p) = 0
    End If
    If Abs(IData(R1n)) < 10 Then
        Rea1 = Abs(AData(m) - _
AData(R1p)) * 2
        IData(R1n) = 0
    End If
'record found r_a in Data2 which
'is an Excel file
    Data2.Refresh
    Data2.Recordset.MoveFirst
    Do While Data2
.Recordset.Fields("USED").Value = "Y"
        Data2.Recordset.MoveNext
    Loop
    If Data2.Recordset.Fields("USED")_
.Value <> "Y" Then
        Data2.Recordset.Edit
        If IData(m) < 1 Then
            Data2
.Recordset.Fields("PEAK").Value = 0_
        Else
            Data2.Recordset
.Fields("PEAK").Value = CSng(VegaZP - _
2.5 * (Log(CDbl(IData(m))) / _
ExpTime / Area) / Log(10#)))
        End If
        If IData(R1n) < 1 Then
            Data2.Recordset.Fields_
("R1SBDPN").Value = 0
        Else
            Data2.Recordset.Fields_
("R1SBDPN").Value = CSng(VegaZP - 2.5_
* (Log(CDbl(IData(R1n))) / _
ExpTime / Area) / Log(10#)))
        End If

```



```

    If IData(R1p) < 1 Then
        Data2.Recordset.Fields_
("R1SBDPP").Value = 0
    Else
        Data2.Recordset.Fields_
("R1SBDPP").Value = CSng(VegaZP -
    2.5 * (Log(CDb1(IData(R1p))) /_
ExpTime / Area) / Log(10#)))
    End If
    Data2.Recordset_
.Fields("USED").Value = "Y"
    Data2.Recordset.Fields_
("SOURCE").Value = Left(File1_
.filename, Len(File1.filename) - 4)
    Data2.Recordset.Fields_
("2REA1").Value = Rea1
    Data2.Recordset.Fields_
("2REA2").Value = Rea2
    Data2.Recordset.Fields_
("PEAKDEG").Value = AData(m)
    Data2.Recordset.Fields_
("M").Value = m
    Data2.Recordset.Fields_

```

```

("R1N1VALUE").Value = AData(R1n)
    Data2.Recordset.Fields_
("R1P1VALUE").Value = AData(R1p)
    Data2.Recordset.Fields_
("NOPIX").Value = p
    Data2.Recordset.Fields_
("PA").Value = PA
    Data2.Recordset.Fields_
("INSTRUMENT").Value = Instrument
    Data2.Recordset.Fields_
("FILTER").Value = Filter
    Data2.Recordset.Fields_
("VEGAZP").Value = VegaZP
    Data2.Recordset.Fields_
("EXPTIME").Value = ExpTime
    Data2.Recordset.Update
Else
    Stop 'got a problem
'with the data file
End If
Data2.Recordset.Close
End Sub

```

-
- [1] N. A. Bahcall et al., *Science* **284**, 1481 (1999).
- [2] E. F. Bell et al., *Mon. Not. R. Astron. Soc.* **343**, 367 (2003).
- [3] F. C. van den Bosch et al., *Astroph. J.* **576**, 21 (2002).
- [4] J. A. Sellwood and A. Kosowsky, in *ASP Conference Series* (2001), vol. 240, p. 311.
- [5] R. Bottema et al., *Astron. Astroph. J.* **393**, 453 (2002), astro-ph/0207469.
- [6] S. McGaugh (1998), astro-ph/9812327.
- [7] E. Battaner and E. Florido (2000), astro-ph/0010475v1.
- [8] Y. Sofue and V. Rubin, *Annual Review of Astronomy and Astrophysics* **39**, 137 (2001).
- [9] M. J. Sempere et al., *Astron. Astroph. J.* **296**, 45 (1995).
- [10] R. Schödel et al., *Nature (London)* **419**, 694 (2002), and references therein.
- [11] C. J. Conselice, M. A. Bershadsky, and A. Jangren, *Astroph. J.* **529**, 886 (2000).
- [12] C. J. Jog, *Astron. Astroph. J.* **391**, 471 (2002).
- [13] J. C. Hodge and M. W. Castelaz (2003), presented at the 202nd meeting of the American Astronomical Society, Session 40.08, astro-ph/0305022.
- [14] F. Donato and P. Salucci (2004), astro-ph/0403206.
- [15] A. M. Ghez et al., *Nature (London)* **407**, 349 (2000).
- [16] L. Ferrarese and D. Merritt (2002), astro-ph/0206222.
- [17] A. M. Ghez (2003), astro-ph/0303151.
- [18] A. M. Ghez et al. (2003), astro-ph/0306130.
- [19] J. Dunning-Davies (2004), astro-ph/0402290.
- [20] N. Prasad and R. S. Bhalerao (2003), astro-ph/0309472.
- [21] Q. D. Wang, E. V. Gotthelf, and C. C. Lang, *Nature (London)* **145**, 148 (2002).
- [22] L. Ferrarese and D. Merritt, *Astroph. J. Lett.* **539**, 9 (2000b).
- [23] K. Gebhardt et al., *Astroph. J. Lett.* **539**, 13 (2000a).
- [24] A. M. Ghez et al., *Astroph. J.* **509**, 678 (1998).
- [25] J. Kormendy and D. Richstone, *Annual Review of Astronomy and Astrophysics* **33**, 581 (1995), and references therein.
- [26] J. Magorrian et al., *Astron. J.* **115**, 2285 (1998), and references therein.
- [27] N. Mouawad et al. (2004), astro-ph/0402338.
- [28] N. Bilic, G. B. Tupper, and R. D. Viollier (2003), astro-ph/0310172.
- [29] J. Binney and M. Merrifield, *Galactic Astronomy* (Princeton, NJ: Princeton University Press, 1998), and references therein.
- [30] A. Königl (2003), astro-ph/0302109.
- [31] F. K. Baganoff et al., *Nature (London)* **413**, 45 (2001).
- [32] F. K. Baganoff, *American Astronomical Society HEAD meeting #35*, session #03.02 (2003a).
- [33] S. Nayakshin and R. Sunyaev (2003), and references therein, astro-ph/0302084.
- [34] J. Zhao et al. (2003), and references therein, astro-ph/0302062.
- [35] D. Merritt and L. Ferrarese (2002), astro-ph/0107134, and references therein.
- [36] K. Gebhardt et al., *Astroph. J. Lett.* **543**, 5 (2000b), astro-ph/0007123.
- [37] L. Ferrarese and D. Merritt (2000a), astro-ph/0006053 v1.
- [38] S. Tremaine et al., *Astroph. J.* **574**, 740 (2002).
- [39] L. Ferrarese, *Astroph. J.* **578**, 90 (2002), astro-ph/0203469.
- [40] C. Shu, H. J. Mo, and S. Mao (2003), astro-ph/0301035.
- [41] J. Silk and M. J. Rees, *Astron. Astroph. J.* **31**, 1 (1998).
- [42] F. K. Baganoff, *Astroph. J.* **591**, 891 (2003b).
- [43] R. Genzel et al., *Nature (London)* **425**, 934 (2003).
- [44] R. J. Mclure and J. S. Dunlop (2002), astro-ph/0201081.
- [45] D. Merritt and L. Ferrarese, *Mon. Not. R. Astron. Soc.*

- 320**, L30 (2001b).
- [46] A. Wandel (2003), astro-ph/0303562.
 - [47] A. W. Graham et al., *Astroph. J.* **563**, 11 (2001).
 - [48] A. W. Graham et al. (2002), astro-ph/0206248.
 - [49] A. W. Graham et al. (2003), and references therein, astro-ph/0306023.
 - [50] H. S. Carslaw and J. C. Jeager, *Conduction of Heat in Solids* (Oxford, NY: Oxford University Press, 2000).
 - [51] A. S. Grove, *Physics and Technology of Semiconductor Devices* (New York, NY: John Wiley and Sons, Inc, 1967), and references therein.
 - [52] C. Will, *Theory and Experiment in Gravitational Physics* (Cambridge, United Kingdom: Cambridge University Press, 1993), and references therein.
 - [53] W. L. Freedman et al., *Astroph. J.* **533**, 47 (2001).
 - [54] S. T. Gottesman, R. D. Davies, and V. C. Reddish, *Mon. Not. R. Astron. Soc.* **133**, 359 (1966).
 - [55] C. Carignan and K. C. Freeman, *Astroph. J.* **294**, 494 (1985).
 - [56] E. Corbelli and P. Salucci, *Mon. Not. R. Astron. Soc.* **311**, 441 (2000).
 - [57] K. G. Begeman, A. H. Broeils, and R. H. Sanders, *Mon. Not. R. Astron. Soc.* **249**, 523 (1991), and references therein.
 - [58] J. A. Fillmore, T. A. Boroson, and A. Dressler, *Astroph. J.* **302**, 208 (1986).
 - [59] A. H. Rots, *Astron. Astroph. J.* **45**, 43 (1975).
 - [60] T. S. van Albada et al., *Astroph. J.* **295**, 305 (1985).
 - [61] E. Moore and S. T. Gottesman, *Mon. Not. R. Astron. Soc.* **294**, 353 (1998).
 - [62] P. C. van der Kruit, *Astroph. J.* **192**, 1 (1974).
 - [63] J. Braine, F. Combes, and W. van Driel, *Astron. Astroph. J.* **280**, 451 (1993).
 - [64] G. Chincarini and R. de Sousa, *Astron. Astroph. J.* **153**, 218 (1985).
 - [65] P. Guhathakurta et al., *Astron. J.* **96**, 851 (1988).
 - [66] A. Bosma, W. M. Gross, and R. J. Allen, *Astron. Astroph. J.* **93**, 106 (1981).
 - [67] P. D. Mannheim and J. Kmetko (2000), and references therein, astro-ph/9602094.
 - [68] L. M. Macri et al., *Astroph. J.* **559**, 243 (2001).
 - [69] R. B. Tully and J. R. Fisher, *Astron. Astroph. J.* **54**, 661 (1977).
 - [70] R. B. Tully and M. J. Pierce, *Astroph. J.* **533**, 44 (2000).
 - [71] J. C. Hodge and M. W. Castela (2003), presented at the 202nd meeting of the American Astronomical Society, Session 40.08, astro-ph/0304030.
 - [72] A. A. Michelson, *Am. J. Of Science* (3) **22**, 20 (1881).
 - [73] M. Consoli and E. Costanzo (2003), astro-ph/0311576.
 - [74] H. Müller et al. (2003), physics/0305117.
 - [75] M. Baes et al., *Mon. Not. R. Astron. Soc.* **341L**, 44 (2003), astro-ph/0303628.
 - [76] T. Padmanabhan and T. R. Choudhury, *Phys. Rev. D* **66**, 081301 (2002).
 - [77] L. Smolin (2004), and references therein, astro-ph/0407213.
 - [78] The NED database is available at: <http://nedwww.ipac.caltech.edu>. The data were obtained from NED on 5 May 2004.
 - [79] Based on photographic data obtained from MAST at <http://archive.stsci.edu/hst/search.php>.
 - [80] The surface brightness was calculated using the VEGAMAG system as described in http://www.stsci.edu/instruments/wfpc2_dhb/wfpc2_ch52.html. The ZEROPOINT for the F450W, F547M, F555W, F606W, F658N, and F702W filters are 22.007, 21.689, 22.571, 22.919, 18.154, and 22.466, respectively.
 - [81] The one-tailed probability that the variances of two arrays are not significantly different as calculated by the Microsoft Excel program.
 - [82] The LEDA database is available at: <http://leda.univ-lyon1.fr>. The data were obtained from LEDA on 5 May 2004.
 - [83] The data and a sample calculation are available in a Microsoft Excel file by e-mail to scjh@citcom.net.